Astronomy & Astrophysics manuscript no. 2005˙HMXBcat.˙4987˙final
February 1, 2008
© ESO 2008

arXiv:0707.0549v1 [astro-ph] 4 Jul 2007

Catalogue of high-mass X-ray binaries in the Galaxy (4th edition) ⋆

Q.Z. Liu[1,2], J. van Paradijs[2], and E.P.J. van den Heuvel[2]

[1] Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, P.R. China
e-mail: qliu@astro.uva.nl
[2] Sterrenkundig Instituut "Anton Pannekoek", Universiteit van Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

Received date/ Accepted date

ABSTRACT

Aims. The aim of this catalogue is to provide the reader with some basic information on the X-ray sources and their counterparts in other wavelength ranges (γ-rays, UV, optical, IR, radio).
Methods. Some sources, however, are only tentatively identified as high-mass X-ray binaries on the basis of their X-ray properties similar to the known high-mass X-ray binaries. Further identification in other wavelength bands is needed to finally determine the nature of these sources. In cases where there is some doubt about the high-mass nature of the X-ray binary this is mentioned. Literature published before 1 October 2005 has, as far as possible, been taken into account.
Results. We present a new edition of the catalogue of high-mass X-ray binaries in the Galaxy. The catalogue contains source name(s), coordinates, finding chart, X-ray luminosity, system parameters, and stellar parameters of the components and other characteristic properties of 114 high-mass X-ray binaries, together with a comprehensive selection of the relevant literature. About 60% of the high-mass X-ray binary candidates are known or suspected Be/X-ray binaries, while 32% are supergiant/X-ray binaries.

Key words. Catalog – X-ray: binaries – Stars: emission-line, Be – Stars: binaries: close

1. Introduction

High-mass X-ray binaries (HMXBs) were among the very first bright X-ray sources detected and optically identified in the 1970s. The systems comprise a compact object orbiting a massive OB class star. The compact object should be either a neutron star (NS) or black hole and is a strong X-ray emitter via the accretion of matter from the OB companion. Thus, cataclysmic variables do not belong to HMXBs (for a catalogue of cataclysmic variables see Ritter & Kolb 2003). Conventionally, HMXBs can be further divided into two subgroups: those in which the primary is a Be star (Be/X-ray binary) and those in which the primary is a supergiant (SG/X-ray binary). One can refer to the book by Lewin & van der Klis (2006) for a full understanding of the various aspects of HMXBs.

The majority of the known high-mass X-ray binaries are Be/X-ray systems (BeXRBs), especially those in the Magellanic Clouds (Liu, van Paradijs & van den Heuvel 2005). Meurs & van den Heuvel (1989) predict 2000–20000 Be/X-ray binaries in the Galaxy. In Be systems, the compact object is a neutron star and is typically in a wide, moderately eccentric orbit, and it spends little time in close proximity to the dense circumstellar disk surrounding the Be companion (Coe 2000; Negueruela 2004). No black hole and Be star system has been found yet (see Zhang, Li & Wang 2004). X-ray outbursts will be expected when the compact object passes through the Be-star disk, accreting from the low-velocity and high-density wind around Be stars, and thus collectively termed Be/X-ray transients. Their X-ray spectra are usually hard. The hard X-ray spectrum, along with the transience, is an important characteristic of the Be/X-ray binaries.

In the second group of HMXB systems, the compact star orbits a supergiant early-type star, deep inside the highly supersonic wind (for a review see Kaper et al. 2004). The X-ray luminosity is either powered by the strong stellar wind of the optical companion or Roche-lobe overflow. In a wind-fed system, accretion from the stellar wind results in a persistent X-ray luminosity of 10^{35}–10^{36} erg s^{-1}, while in a Roche-lobe overflow system, matter flows via the inner Lagrangian point to an accretion disc. A much higher X-ray luminosity ($\sim 10^{38}$ erg s^{-1}) is produced.

Five years after the publication of the previous (3rd) edition (Liu, van Paradijs & van den Heuvel 2000), the amount of new literature and the number of new objects to be included have again grown so much that it seems worthwhile to concentrate the information of HMXBs in the Magellanic Clouds and in the Galaxy separately. In a previous paper we published a catalogue of 128 HMXBs in the Magellanic Clouds (Liu, van Paradijs & van den Heuvel 2005), and here we present a catalogue of HMXBs in the Galaxy. We briefly recall some of the developments that, over the past five years, have had (and still have) a major impact on this catalogue.

Due to the much increased sensitivity and spatial resolution achievable with the Chandra and the XMM-Newton X-ray observatories, as well as with the Hubble Space Telescope and large ground-based radio telescopes, more accurate positions of X-ray binaries have been determined, resulting in the unambiguous discovery of the optical and/or IR counterpart to some X-ray sources. Moreover, the number of HMXBs in external galaxies is also rapidly increasing, e.g., the X-ray binaries in M81 (Swartz et al. 2003) and the Antennae galaxies (Zezas et al. 2002). Most of the ultra-luminous X-ray sources in starburst and spiral galaxies (Liu & Mirabel 2005) are believed to be HMXBs with a black hole, specifically the B0 Ib star to NGC 5204 X-1 (Liu et al.

Send offprint requests to: Q.Z. Liu

⋆ Table 1 and the references are only available in electronically form at the CDS via anonymous ftp (130.79.128.5) or via http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/???/????

2004), the O9I star to M 33 X-7 (Pietsch et al. 2004) and the mid-B supergiant to M101 ULX-1 (Kuntz et al. 2005). It has been proposed that the collective X-ray luminosity of high-mass X-ray binaries can be used as an indicator of the star-formation rate for the host galaxy (Grimm, Gilfanov & Sunyaev 2003).

Since its launch in 2002, INTEGRAL has been revealing hard X-ray sources that were not easily detected in earlier soft X-ray (typically ≤10 keV) observations. Hard X-rays are not easily absorbed by matter and thus are highly penetrating. Such radiation is, therefore, ideal for probing high-energy emitting sources in dense regions. INTEGRAL has been performing a regular survey of the Galactic plane and a deep exposure of the Galactic Center as part of its Core Program (Winkler et al. 2003). A group of hard X-ray sources emitting in the hard X-ray and soft γ-ray regions have been discovered in the course of the INTEGRAL observations, which are highly absorbed, i.e., with column densities higher than about 10^{23} cm^{-2}. The X-ray, as well as the optical/IR, properties of these sources and their location in the sky suggest that they may belong to the class of high-mass X-ray binaries, some of them possibly long-period X-ray pulsars. The donors in these binaries are most probably giant or supergiant stars (see Kuulkers 2005). For the details of all INTEGRAL sources please refer to the web page of Jerome Rodriguez (http://isdc.unige.ch/~rodrigue/html/igrsources.html).

In recent years there has been growing evidence that there is a class of X-ray binaries referred to as fast X-ray transients, characterized with X-ray outburst durations on the order of a few hours and peak fluxes on the order of 10^{-9} erg cm^{-2} s^{-1}(2–10 keV). They are not readily identified with other similarly active X-ray sources: magnetically active nearby stars (i.e., DY Dra, RS CVn, or pre-main sequence stars) or superbursters (e.g., Kuulkers 2004). These sources lie in the vicinity of the Galactic Center, a region extensively monitored by INTEGRAL and other satellites. The high fluxes and the lack of nearby counterparts suggest high luminosities that would indicate an X-ray binary origin. Several of these fast transients are associated with OB supergiants (Coe et al. 1996; Halpern et al. 2004; Negueruela et al. 2006; Smith et al. 2006; Pellizza et al. 2006), so they are suggested to be a new class of high-mass X-ray binaries (Negueruela et al. 2005).

The aim of our catalogue is to provide some basic information on the X-ray sources and their counterparts, as well as the binary properties of the system in question, and easy access to the recent literature. No attempt has been made to compile complete reference lists. Much effort has been made to avoid errors and to keep the information up to date. Nevertheless, the authors are well aware that this edition too may contain errors and may be incomplete.

2. Description of the table

Table 1 lists the 114 HMXB candidates in the Galaxy. The format of Table 1 is similar to the previous edition and almost the same with the HMXB catalogue in the Magellanic Clouds (Liu, van Paradijs & van den Heuvel 2000; 2005), of which the present catalogue is meant to be an update. If there is any doubt about an entry, a question mark follows the item. In the table the sources are ordered according to the right ascension of sources; part of the (mainly numerical) information on a source is arranged in six columns, below which additional information is provided for each source in the form of key words with reference numbers [in square brackets]. The columns have been arranged as follows.

In Column 1 the first line contains the source name, with rough information on its sky location, according to the conventional source nomenclature of space experiments in which the source was detected, hhmm±ddd or hhmm.m±ddmm. Here hh, mm, and ss indicate the hours, minutes and seconds of right ascension, ddd the declination in units of 0.1 degree (in a small number of cases, the coordinates shown in the name are given with more, or fewer, digits). The prefix J indicates a name based on J2000 coordinates. Otherwise, 1950 coordinates were used in the name. An alternative source name is given in the second line. In the third line of Column 1, the source types are indicated with a letter code, as follows:

- P: X-ray pulsar (66);
- T: transient X-ray source (62);
- E: eclipsing system (9);
- R: radio emitting HMXBs (9);
- C: cyclotron resonance scattering feature at X-ray spectrum (16);
- U: ultra-soft X-ray spectrum (1). These sources include black-hole candidates (BHC); some 'extreme ultra-soft' (EUS) source may be a white dwarf (WD) on whose surface steady nuclear burning takes place.

Column 2 contains the right ascension (RA) and declination (DEC) of the source for equinox J2000.0 in the first two lines. RA is given as hhmmss.s to an accuracy of 0.1 s, DEC is given in ° ' ", to an accuracy of 1" (in a small number of cases, the coordinates are given with more, or fewer, digits). The third line gives the galactic longitude and latitude to an accuracy of 0.1° (except for sources close to the galactic center, where these coordinates are given to 0.01°). A reference for the source position is given below the columnar information under *'pos.'*. In the parentheses following the *'pos.'*, we provide some information on the type of observation from which the source position has been derived. The following abbreviations are used: o, optical; x, X-ray; r, radio; IR, infrared. Following the type of observation, we give an indication of the accuracy of this position, in the form of equivalent (90 percent confidence level) error radii, but in several cases this can only be considered an approximation (e.g. when the error box is not circular). When no accuracy is quoted, it is about one arcsecond or better.

The first and second lines of Column 3 give the names of the optical counterpart to an X-ray source. The third line contains a reference to a finding chart. An asterisk followed by a number or letter refers to the star number used in the finding chart; "star" refers to a star in the finding chart that has not been assigned a number or letter. Many optical counterparts have been indicated with a variable-star name, as given in the *General Catalogue of Variable Stars* and in recent name lists of variable stars as published regularly in the *IAU Information Bulletin on Variable Stars*, or a number in a well-known catalogue (e.g., HD, SAO, GSC, 2MASS).

The fourth column contains some photometric information on the optical counterpart. In the first line, the apparent visual magnitude, V, and the color indices $B-V$ and $U-B$ are listed. The second line contains the spectral type of the optical counterpart and an estimate of the interstellar reddening, E_{B-V}.

The fifth column lists the near-infrared magnitudes at J (1.25 microns), H (1.65 microns), and Ks (2.17 microns), detected with the Two Micron All Sky Survey (2MASS)(Skrutskie et al. 2006).

In Column 6, the maximum X-ray flux, or the range of observed X-ray fluxes (2–10 keV, unless otherwise indicated), is

given in units of

$$1\mu Jy = 10^{-29}\ erg\ cm^{-2}\ s^{-1}Hz^{-1} = 2.4\times10^{-12}\ erg\ cm^{-2}\ s^{-1}\ keV^{-1}.$$

The first line in Column 7 gives the orbital period in days. The second line contains the pulse period for X-ray pulsars, in seconds. The third line contains a reference in which the orbital and/or pulse periods were detected.

3. Conclusions and remarks

The current version of this catalogue provides tabulated data and references for 114 objects, including 35 newly discovered HMXBs (2 previously listed in our low-mass X-ray binary catalogue), as well as 79 "old" ones listed in the previous catalogue. Compared with the 3rd edition, the number of HMXBs in the Galaxy listed has increased by ∼43%. Among the 114 HMXB candidates, we find 39(13) confirmed(suspected) Be/X-ray binaries (another 4 dubious Be/X-ray binaries, probably a white dwarf with a Be companion), 18(11) SG/X-ray binaries, and 6 other sources with peculiar features (CI Cam, IGR J16318-4848, Cyg X-3, LS 5039, V4641 Sgr, and SS 433). The remaining 23 systems are candidates of HMXBs. No detailed information on optical companions is known.

We would like to make some remarks on several individual sources. Both KS 1947+300 and GRO J1948+32 have been listed in the 3rd edition. Swank & Morgan (2000), however, found that the transient X-ray source KS 1947+300 has pulsations that are identical to the pulsar GRO J1948+32, and concluded that they are the same source. Also, the ROSAT source, RX J1037.5−5647, is the only one in the Uhuru error boxes 4U 1036−56 (Motch et al. 1997), so probably having the same origin.

MXB 0656−072 was described as a soft X-ray transient during the 1975 outburst (Clark 1975) and previously catalogued as a low-mass X-ray binary. However, observations with the PCA on board the RXTE satellite in October 2003 indicated that the source is actually a hard X-ray pulsar with a spin period of 160.7 s (Morgan, Remillard & Swank 2003). Moreover, the position of the ROSAT counterpart, RX J065817.7−071228, is consistent with a variable-reddened O9.7Ve spectral type Be star, suggesting a Be X-ray binary nature (Pakull, Motch & Negueruela 2003). Another source, SAX J1819.3−2525/V4641 Sgr, was previously classified as a low-mass X-ray binary. Although the accretion processes in this system are typical of those seen in LMXBs, recent spectroscopic observations yielded a mass of the compact star in the range 5.49–8.14 $M_\odot$, and the optical companion was found to be a B9III star (Orosz et al. 2001). Chaty et al. (2003) also showed that the stellar type of the companion star was early type between B3 and A2 and that the binary was between intermediate-mass X-ray binary and HMXB. Therefore, this system is more consistent with a HMXB than a LMXB.

We also wish to emphasize that some sources listed in this catalogue are still uncertain. They should be considered with caution, in view of all the further work needed. Some sources are tentatively classified as massive X-ray binaries due to their X-ray properties, for instance, a hard X-ray transient, a fast X-ray transient, or an extremely absorbed X-ray source. No counterpart at other bands has been found or no pulse period has been detected, or both. Also, the compact object in some weak or soft X-ray sources may be a white dwarf instead of a neutron star, e.g., 1H 0556+286, 1H 0749-600, 1H 1249-637, and 1H 1253-761 (Torrejon & Orr 2001 and see Table 1). If this is the case, they should be excluded from this catalogue.

IGR J12349−6434 is tentatively identified as a HMXB candidate, primarily due to an early-type star, RT Cru, in the error of the X-ray source (Masetti et al. 2005), while XTE J1859+083 is included simply due to the detection of X-ray pulsations (Marshall et al. 1999). Highly absorbed hard X-ray sources are suggested as being high-mass X-ray binaries (see Kuulkers 2005). Two hard X-ray sources, IGR J16418−4532 and IGR J16493−4348, with column densities of about 10^{23} cm^{-2} (Walter et al. 2004; Markwardt et al. 2005), are also included in this catalogue. Fast X-ray transients are probably another new class of high-mass X-ray binaries with a supergiant companion (Smith et al. 2006; Negueruela et al. 2006). The two fast transients, AX J1749.1−2733 (Sakano et al. 2002, see in't Zand 2005) and XTE J1901+014 (Remillard & Smith 2002), together with 8 other members (XTE J1739-302, IGR J17544-2619, IGR J16465-4507, AX 1845.0-0433, AX J1841.0-0536, SAX J1818.6-1703, IGR J16479-4514, and IGR J11215-5952), are included in this edition.

Finally, the following four sources are no longer listed in our catalogue. Now it is quite clear that they do not belong to HMXBs.

1E 1024.0−5732/Wack 2134 was discovered with the Einstein Observatory. Caraveo et al. (1989) suggested that a highly reddened O5 star was the most likely optical counterpart, and therefore proposed the system was a HMXB. The source was listed in the previous catalogue. Optical and X-ray data, however, favor the system containing a Wolf-Rayet star and an O-type star (Mereghetti et al. 1994; Reig 1999). The origin of the X-rays from this source is explained by the colliding-wind binary model (Reig 1999).

1E 1048.1−5937 and GS 1845−03 were listed in the previous catalogue. GS 1845−03 is, however, probably the same as the ASCA source, AX J1845.0−0258. Both the compact star of 1E 1048.1−5937 and AX J1845.0−0258 (Torii et al. 1998; Gotthelf & Vasisht 1998) very likely belong to the "anomalous" X-ray pulsars, which are now thought to be related to the magnetars. Mereghetti et al. (1998) argued that if 1E 1048.1−5937 has a companion this star must be either a white dwarf or a helium-burning star instead of a main-sequence companion.

The peculiar galactic X-ray source, 4U 1954+319, has been listed in the previous editions as an HMXB. Masetti et al. (2006), however, found that the suspected field M-type giant star is indeed the counterpart of the X-ray source, based on the Chandra Observation. They suggest that 4U 1954+319 is a wide-orbit LMXB, most likely a neutron star, accreting from the wind of an M-type giant.

V669 Cep used to be identified as the optical counterpart to the hard X-ray source, 1H 2214+589. However, in't Zand et al. (2000) identify another Be star as the optical counterpart to its BeppoSAX counterpart, SAX J2239.3+6116. Instead, V669 Cep is identified as the optical counterpart to RX J2226.6+6113 (Halpern et al. 2001). Both Hang, Liu & Xia (1999) and Halpern et al. (2001) classify V669 Cep as a Herbig Ae/Be star, while Miroshnichenko et al. (2002) suggest it as a binary, consisting of a hot, low luminosity B4–B6 star and a cool companion, most likely a late-type giant. We have excluded this source in the 4th edition, since in both cases RX J2226.6+6113/V669 Cep is unlikely to be an HMXB, although it is still a HMXB in SIMBAD.

Noted added in proof: After we submitted the paper, we noticed that Swift J1626.6-5156 (Negueruela et al. 2006, ATel 739), XTE J1716-389 (Cornelisse et al. 2006, MNRAS, 366, 918), IGR J16207-5129, and AX J1700.2-4220 (Palazzi, et al. 2006, ATel 783) are likely candidates for high-mass X-ray binaries.

Acknowledgements. We wish to thank Marc Ribó, Sylvain Chaty and Ignacio Negueruela for carefully reading the manuscript and for their useful comments. We also thank Leonardo Pellizza and Sylvain Chaty for providing us with information on IGR J17544−2619 prior to publication. Finally, we are very grateful to the referees, Malcolm Coe and Sebastien Derriere, for their careful and thorough reading of this paper.

This research has made use of the SIMBAD data base operated at the CDS, Strasbourg, France, and NASA's Astrophysics Data System (ADS). This publication also makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

QZL is partially supported by the Royal Science Foundation of The Netherlands, the National Natural Science Foundation of China under Grants 10173026 and 10433030, and the Major State Basic Research Development Program of China (973 Program) under Grant G1999075405.

Table 1. High-mass X-ray binaries in the Galaxy

Name(s) type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
IGR J00370+6122 1RXS J003709.6+612131	00 37 10 +61 21 35 121.2, −1.5	BD +60 73 LS I +61 161	9.65, 0.56, B0.5II–III, 0.75 [640]	8.389 8.265 8.166	8 (2–28 keV) [165]	15.665 [165]
Pos. (o): [640]; distance of 3.0 kpc: [640]; variable hard X-ray source: [165]; no radio emission detected: [75].						
2S 0053+604 3A 0053+604 R	00 56 42.5 +60 43 00 123.6, −2.1	γ Cas [175]	1.6–3.0, −0.15, −1.08 B0.5 Ve, 0.05 [528]	2.039 1.993 1.756	5–11 [64]	203.59 [279]
Pos. (o): [64]; X-ray spectrum: [220, 545, 588]; UV observations: [169, 275, 286, 759]; variable Be star: [469, 810]; radial-velocity variations: [358]; $v_r sini \sim$ 300–500 km s^{-1}: [312, 318, 528]; system velocity: [795]; radio/mm obs.: [178, 817]; polarimetry: [107]; optical interferometry Hα envelope: [542, 763]; non-radial pulsations: [855]; wind structure: [816]; a WD sys.: [263]; multiwavelength (mwl) obs.: [305, 716]; ISO spe. and envelope structure: [304]; secondary mass of $\sim 1 M_\odot$: [279]; $e = 0.26$: [279]; X-ray and optical variations: [664]; multi-component of X-ray emission: [717]; distance of 0.19 kpc: [598].						
2S 0114+650 P, C?	01 18 02.7 +65 17 30 125.7, +2.6	V662 Cas LS I +65°010 [175]	11.0, 1.2, 0.1 B0.5 Ib, 1.4 [2, 148]	8.597 8.296 8.107	4 [64]	11.6 10008 [148, 210]
Pos. (o): [598]; X rays: [210, 211, 391, 486]; opt. spectra: [2, 148, 434]; $v_r sini \sim$ 45 km s^{-1}: [2]; opt. polarimetry: [41]; magnetar origin: [430]; astrophys. parameters: [633]; opt. photometry: [758]; superorbital period of 30.7 d: [194]; distance of 7.2 kpc: [633]; spin-up: [55]; likely a cyclotron resonance scattering feature at 22 keV with one harmonic (implying a magnetic field of 2.5×10^{12} G) [55], but not detected with SAX: [486].						
4U 0115+634 4U 0115+63 T, P, C	01 18 31.9 +63 44 24 125.6, +1.0	V635 Cas [361]	14.5–16.3, 1.4, 0.3 B0.2Ve, 1.5 [507, 554, 784]	11.622 11.023 10.629	<2–350 [64]	24.3 3.61 [547, 625]
Pos. (o): [361]; X-ray outbursts: [614, 659, 675, 750, 776]; recurrence time 3 yr (?): [831]; rapid X-ray variation: [77]; X-ray spectrum: [659, 830]; X-ray cyclotron lines: [282, 549, 750]; opt. pre-outburst behavior: [407, 507]; long-term opt. phot.: [507]; opt. spectra: [317, 784]; $v_r sini \sim 365 km/s$: [317]; mwl obs. during an outburst: [557]; distance of 7–8 kpc: [554]; $e = 0.34$: [625]; QPOs during outburst: [110, 721].						
IGR J01363+6610 T	01 35 50 66 12 40 127.4, 3.7	EM* GGR 212 [640]	13.29, 1.39, B1Ve, 1.61 [640]	10.039 9.565 9.121	9 (17–45 keV) [251]	
Pos. (o): [640]; distance of 2.0 kpc: [640]; discovered by INTEGRAL: [251]; very strong Hα emission with EW of −50Å: [640]; lack of radio emission: [606].						
RX J0146.9+6121 	01 47 00.2 +61 21 23.7 129.5, −0.8	LS I +61°235 V831 Cas [278]	11.45, 0.62, −0.41 B1Ve, 0.87 [116, 634]	9.899 9.700 9.486	1–3 (0.1–2.4 keV) [538]	 1404.2 [266]
Pos. (o): [602]; in open cluster NGC 663: [687]; $v_r sini \sim$ 200 km s^{-1}: [634]; X-ray obs.: [510]; X-ray outburst: [266]; physical parameters and V/R variability: [634]; long-term optical and IR variability: [636]; distance of 2.3 kpc: [116].						
IGR J01583+6713 T	01 58 18.44 67 13 23.5 129.4, 5.2	 	 Be, [270]	11.481 11.033 10.601	14 (20–40 keV) [729]	
Pos. (o): [270]; hard X-ray transient, not detected in the 40-80 keV band: [729]; 7.7' from 1RXS J015918.4+671807: [729]; USNO-B1.0 POSS II magnitudes of B=14.98, R=13.24, I=12.12; strong Hα and weak Hβ emission lines, typical features of a Be/X-ray binary: [270]; highly absorbed, with N_H approximately 10^{23} cm^{-2}: [380].						

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
1E 0236.6+6100	02 40 31.7	LS I +61°303	10.7, 0.8, −0.3	8.621	0.2	26.496
	+61 13 46	V615 Cas	B0Ve, 0.75	8.228	(0.2–5 keV)	
R	135.7, +1.1	[278]	[179, 316, 726]	7.917	[43]	[755, 254]

Pos. (o,r): [256]; X-ray source: [43]; two-peak accretion/ejection model: [757, 478, 496]; radio outbursts: [755, 756]; long-term radio obs.: [629]; four-year modulation radio outbursts: [184, 579, 254]; fast precessing relativistic radio jet: [495]; microquasar (μQ): [494]; VLBI radio jet: [493, 757]; optical light curve: [506, 578]; opt. spectra: [437, 316, 392]; orbital solution: [316, 478, 79, 254]; $v_r sini \sim 113$ km s^{-1}: [79]; system velocity: [795]; long-term optical obs.: [256, 438, 437]; IR obs.: [156, 311]; distance about 2.4 kpc: [217, 726]; UV obs.: [306, 315, 316, 463]; wind structure: [816]; in the error of 3EG J0241+6103: [388]; simultaneous X-ray/radio obs.: [280]; low-mass BHC: [621]; mwl obs.: [725]; orbital period: [580, 581, 856, 492, 418, 254]; eccentricity of 0.72: [478, 79]; ejected from cluster IC 1805: [522].

V 0332+53	03 34 59.9	BQ Cam	15.1–15.4, 1.6–2.3,	11.817	<0.5–1250	34.25
EXO 0331+530	+53 10 24		O8.5Ve, 1.9	11.206		4.4
T, P, C	146.1, −2.2	[389]	[137, 302, 552]	10.744	[730, 761]	[547, 730, 832]

Pos. (o): [389]; X-ray outbursts: [730, 761, 832]; X-ray obs.: [461]; rapid X-ray variability: [35]; X-ray cyclotron line: [460]; multiple cyclotron lines: [108]; X-ray pulse phase sp.: [783]; QPO: [747, 622]; optical spectrum: [354, 737]; $v_r sini \sim 150$ km s^{-1}: [137]; system velocity: [795]; IR obs.: [113]; wind structure: [816]; tilted equatorial plane: [553]; distance of 7 kpc: [553]; eccentricity of 0.37: [858, 730].

4U 0352+309	03 55 23.0	X Per	6.0–6.6, 0.29, −0.82	6.149	<9–37	250.3
	+31 02 59		B0Ve, 0.4	6.073		835
P, C	163.1, −17.1	[65]	[319, 451, 528, 710]	5.920	[64]	[162, 547]

Pos. (o): [64]; X-rays: [660]; X-ray sp.: [830]; long-term optical record: [435, 532]; disappearance emission lines: [435, 566]; system velocity: [795]; $v_r sini \sim 250$–400 km s^{-1}: [314, 528, 827]; UV obs.: [40, 275]; stellar parameters: [189]; opt. polarimetry: [409]; envelope structure: [760]; photometry and polarimetry obs. during disc loss: [666]; fundamental parameters: [451]; mwl study: [665]; cyclotron line ($B = 3.3 \times 10^{12}$ G): [109]; orbital parameters ($e = 0.11$): [162]; distance of 0.7–1.3 kpc: [189, 451, 666, 760].

XTE J0421+560	04 19 42.2	CI Cam	9.25–11.83, 1.0, −0.4	7.502	2000	19.41
	55 59 59		sgB[e], 0.8	5.741		
T, R	149.2, +4.1	[176]	[104, 105, 327]	4.323	[37, 712]	[28]

Pos. (o): [394]; rapid X-ray outbursts: [221, 712]; mwl obs.: [37]; optical outbursts: [221, 663]; B[e]: [37]; sgB[e] star: [104, 663, 327]; IR spe.: [104]; opt./IR/radio obs.: [105]; long-term UBVRIJHK obs.: [39]; a WD: [571, 221, 346, 28]; likely a BHC: [37, 663]; distance 1–>5 kpc: [37, 104, 519, 346, 523, 663]; E(B−V) between 0.65 and 1.54: [104, 663, 571]; orbital parameters (e=0.62): [28]; B4 III–V: [28]; unusually strong FeII lines: [176]; two-component wind: [663]; radio obs. (failed jet?): [519]; flare peaked successively from X-ray, optical to radio: [221]; rapid changed X-ray N_H: [37, 589].

RX J0440.9+4431	04 40 59.3	LS V +44 17	10.78, 0.61, −0.36	9.500	1	
	44 31 49	BSD 24-491	B0.2Ve, 0.65	9.317		202.5
P	159.8, -1.3	[145]	[539, 641]	9.182	[632]	[632]

Pos. (o): [300]; distance 3.3 kpc: [539, 641]; optical spe.: [641]; Hα EW decreasing with IR magnitude fainter: [641]; *vsini*=235 km s^{-1}: [641].

EXO 051910+3737.7	05 22 35.2	V420 Aur	7.51, 0.14, −0.86	6.696	1
1H 0521+373	+37 40 34	HD 34921	B0 IVpe, 0.42	6.507	
	170.0, +0.7	[719]	[74, 604, 676]	6.230	[215, 850]

Pos. (o): [719]; EXO 051910+3737.7 not the same X-ray src of 1H 0521+373 in SIMBAD; hard X-ray spectrum: [604]; UV/opt./IR obs.: [604, 676]; IR spe.: [104]; unclB[e] star: [104].

1A 0535+262	05 38 54.6	V725 Tau	8.9–9.6, 0.45–0.62, −0.54	8.368	<3–2800	111
H 0535+262	+26 18 57	HD 245770	O9.7IIIe, 0.8	8.272		104
T, P, C	181.4, −2.6	[429]	[228, 465, 726, 734]	8.157	[64]	[547, 616]

Pos. (o): [64]; long-term X-ray obs.: [616]; hard X-ray obs.: [115, 655, 699]; long-term optical record: [38, 734, 452]; opt. spectr.: [3, 320, 324]; $v_r sini \sim 300$ km s^{-1}: [320]; system velocity: [795]; mwl obs.: [164, 356]; X-ray/opt. relation: [454]; UV obs.: [164, 594, 853]; accretion model: [537]; wind structure: [816]; disc loss & renewal: [267]; IR spe. obs.: [103]; long-term opt./UV spe.: [102]; cyclotron lines: [260, 404]; QPO during a giant outburst: [208]; orbital elements (e=0.47): [208]; distance of 2 kpc [726] and 2.9 kpc in [452]; X-ray pulsations during disc loss phase: [558]; variable γ-ray emission: [672].

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
1H 0556+286	05 55 55.1 +28 47 06 181.3, +1.9	HD 249179	9.2, , B5ne, [608, 778]	9.366 9.238 9.068	1.1 [850]	
Pos. (o): [778]; Be star: [810]; distance of 0.83 kpc: [54]; likely not an X-ray binary: [771].						
IGR J06074+2205 T	06 07 26.6 22 05 483		 Be, [272]	10.49 10.19 9.96	7 (3–10 keV) [88]	
Pos. (o): [272]; discovery: [88]; B = 13.3, R = 12.1: [272]; opt. emission line, a Be/X-ray binary: [272]; the radio source unrelated: [272].						
SAX J0635.2+0533 SAX J0635+0533 P	06 35 18.29 05 33 06.3 206.1, −1.0	 [363]	12.83, +0.98, B2V–B1IIIe, 1.2 [363]		0.55 [363]	11.2 0.0338 [151, 364]
Pos. (o): [363]; opt. emission line: [363]; within the error of 2EG J0635+0521: [363]; distance ∼ 2.5–5 kpc: [363]; eccentr. of 0.29: [364].						
1WGA J0648.0−4419 RX J0648.1−4419 P	06 48 04.6 −44 18 58.4 253.7, −19.1	HD 49798	8.27, −0.24, −1.18 sdO6, [359]	8.862 9.005 9.072	0.03 [347]	1.55 13.1789 [347, 762]
Pos. (o): [598]; distance 650 pc: [347]; very soft X-ray: [347]; UV sp.: [71]; likely a NS: [347]; a WD system(?): [45]; eccentricity of 0.03: [762].						
XTE J0658−073 MXB 0656−072 T, P, C	06 58 17.30 −07 12 35.3 220.1, −1.8	[M81] I-33	12.05–12.38, 1.02–0.86, O9.7Ve, [576]	9.664 9.332 9.013	67 (2–12 keV) [534]	 160.7 [534]
Pos. (o): [576]; also RX J065817.7−071228 and optical ctpt. GSC2 S100221212: [576]; typical of accretion-powered pulsars: [534]; hard sp. with α=1.09: [285]; previously "soft X-ray transient": [97]; strong cyclotron line at ∼35keV (B about 4×10^{12} G): [285]; variable reddened: [576]; extremely narrow emission lines (HI, HeI and low-ionization metals): [576].						
3A 0726−260 4U 0728−25 T, P	07 28 53.6 −26 06 29 240.3, −4.1	V441 Pup [728]	11.6, 0.3, −0.6 O8–9Ve, 0.6 [556, 728]	10.366 10.093 9.832	1.2–4.7 [815, 850]	34.5 103.2 [138]
Pos. (o): [728]; distance 4.6 or 6.1 kpc: [131, 556]; X-ray flare: [728]; opt./IR obs.: [556]; orbital and spin periods: [138].						
1H 0739−529	07 47 23.6 −53 19 57 266.3, −13.7	HD 63666 SAO 235515 [719]	7.62, 0.02, −0.24 B7 IV–Ve, [456, 564, 778]	7.403 7.406 7.363	0.7 [850]	
Pos. (o): [778]; parallax and proper motion: [94]; distance of 0.52 kpc: [94].						
1H 0749−600	07 50 15.77 −61 05 58.2 272.8, −16.5	HD 65663 SAO 250018 [719]	6.73, 0.05, −0.25 B8 IIIe, 0.09 [152, 153, 778]	6.542 6.571 6.534	0.7 [850]	
Pos. (IR): [778]; Be star: [810]; in open cluster NGC 2516: [152]; parallax and proper motion: [94]; distance of 0.4 kpc: [94]; likely not an X-ray binary: [771].						
RX J0812.4−3114 T, P	08 12 28.4 −31 14 21 249.6, +1.6	LS 992 V572 Pup [539]	12.48, 0.38, −0.29 B0.2IVe, 0.65 [637]	11.335 11.126 10.909	0.6 [539]	81.3 31.8851 [133, 631]
Pos. (IR): [539]; distance about 9 kpc: [539, 637]; sharp dip in the pulse profile: [631]; Balmer lines from absorption to emission and back again to absorption: [637]; IR mag. varied by more than 0.8 mag: [637]; *vsini* = 240 km s^{-1}: [637].						

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
GS 0834−430	08 35 55.4	*D	20.4, ,		30–300	105.8
	43 11 11.9		B0–2 III–Ve, 4.0			12.3
T, P	177.7, 36.7	[349]	[349]		[249]	[249, 836]

Pos. (o): [349]; sequence X-ray outbursts: [836]; recurrent transient: [416]; X-ray obs.: [11]; X-ray/opt. obs.: [36]; eccentricity $0.1 < e < 0.17$: [836]; strong Hα emission: [349]; spectral-type uncertain: [349]; distance 3–5 kpc: [349]; southern of two neighboring sources: [123, 249]; IR/opt obs.: [349].

4U 0900−40	09 02 06.9	HD 77581	6.9, 0.47, −0.51	5.833	2–1100	8.96
Vela X-1	−40 33 17	GP Vel	B0.5 Ib, 0.7	5.705		283
P, E, C	263.1, +3.9	[213]	[182, 564]	5.596	[64]	[161, 547, 786]

Pos. (o): [64]; X-ray sp.: [548, 690, 830]; cyclotron line (B of 2.7×10^{12} G): [405, 412]; pulse timing & X-ray orbit: [62, 63, 160, 161, 627, 775, 786]; X-ray eclipses: [213, 781]; high-resolution X-ray spectrum: [694, 241]; limits on apsidal motion: [159]; rapid X-ray variability: [35]; orbital variation X-ray absorption: [264]; energy dependence pulse profile: [626]; opt. light curve: [766]; long-term cycles (?): [614, 766]; opt. radial-velocity curve: [793, 797]; system velocity: [795]; $v_r sini \sim$ 90–130 km s^{-1}: [514, 766, 833]; UV obs.: [182, 365, 502, 594, 682, 794]; high-resolution IUE obs.: [733]; opt. polarimetry: [172]; IR obs.: [238, 326]; bow shock around the sys.: [369]; polarization: [849]; distance of 1.9 kpc [682]; runaway system from OB association Vel OB1: [800]; $1.86< M_x < 2.27 M_\odot$: [29, 623]; mass of GP Vel of $27.9 M_\odot$: [623]; radio emission not detected: [197].

GRO J1008−57	10 09 46.9	star	15.27, −1.66,	10.943	1200	135
	−58 17 35.5		B0e, 1.9–2.0	10.272		93.5
T, P	283.0, −1.8	[117]	[117, 657]	9.718	[738]	[706, 738]

Pos. (o): [117]; X rays: [706]; IR and optical spectral obs.: [117]; 260 days orbital period: [206]; e=0.66 and orbital period of 248 d: [ref. 568]; distance 5 kpc: [117]; large IR excess: [117].

RX J1037.5−5647	10 37 35.2	LS 1698	11.3, 0.9,		3.3	
4U 1036−56	−56 47 59		B0V–IIIe, 0.75			862
P, T	285.4, +1.5	[539]	[539]		[632, 815]	[632]

Pos. (x 21"): [539]; X-ray flare 1974 Nov.: [815]; distance 5.0 kpc: [539]; 2 2MASS sources in the error.

1A 1118−615	11 20 57.2	Hen 3-640	12.1, 0.96, −0.30	9.563	0.1–70	
	−61 55 00	WRAY 15-793	O9.5Ve, 1.2	9.071		405
P, T	292.5, −0.9	[355]	[355]	8.587	[64, 536]	[547]

Pos. (o): [64]; X-ray transient: [186, 352]; opt. phm. & spe.: [355]; sys. velocity: [795]; $v_r sini \sim$ 300 km s^{-1}: [355]; X-ray/opt. obs.: [536]; opt. spe. after X-ray outburst: [605]; mwl study: [118]; UV obs.: [112]; distance of 5 kpc: [355].

4U 1119−603	11 21 15.1	V779 Cen	13.3, 1.07, −0.04	10.736	10–312	2.09
Cen X-3	−60 37 25.5		O6.5 II–III, 1.4	10.311		4.84
P, E, C	292.1, +0.3	[70, 408]	[321, 408, 656]	10.093	[64]	[374, 547]

Pos. (x 0.6"): [328]; X-ray pulsations: [227, 775]; period decay: [187, 374]; X-ray spectrum: [17, 445, 830, 847]; long-term X-ray history: [261, 345, 611, 617, 693]; X-ray eclipses: [692]; X-ray obs.: [307, 544, 550]; aperiodic X-ray variability: [35]; QPO: [746]; cyclotron line at ∼30 keV (B of $\sim 3.5\times10^{12}$ G): [72]; opt. light curve: [766, 798]; opt. radial-velocity curve: [321, 541, 573]; mass of the NS: [15]; $v_r sini \sim$ 250 km s^{-1}: [321]; system velocity: [795]; IR obs.: [238]; structure companion star: [101]; distance of 8–10 kpc: [321, 408]; outburst of GeV γ ray and GeV pulsation: [804].

IGR J11215−5952	11 21 46.81	HD 306414	9.98, 0.59, −0.45	8.55	42
	−59 51 47.9		B1Ia, 0.78	8.34	(20–60 keV)
T	291.9, 1.1		[387, 807]	8.19	[444]

Pos. (o): [561]; about 45' from Cen X-3: [444]; three brighter IR sources in the error: [561]; distance ∼8 kpc: [561].

IGR J11435−6109	11 44 10.7		11.486	10	52.46
	−61 07 02	B3e	11.236	(18–45 keV)	161.76
T, P	294.9, 0.7	[135]	11.107	[252]	[335, 744]

Pos. (o): [135]; B = 13.17: [135]; X-ray variable: [744]; ≠1RXS J114358.1−610736: [135]; optical ctpt. USNO-B1.0 0288−0337948: [135].

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
2S 1145−619	11 48 00.0	V801 Cen	9.3, 0.18, −0.81	8.677	4–1000	187.5
2E 1145.5-6155	−62 12 25	HD 102567	B0.2IIIe, 0.35	8.577		292.4
P, T	295.6, −0.2	[65]	[568, 564, 816]	8.403	[64]	[547, 819]
Pos. (o): [64]; X-ray spectrum: [830]; X-ray obs.: [508]; long-term X-ray history: [617, 819]; opt. spectrum: [828]; $v_r sini \sim 270 km/s$: [275]; system velocity: [795]; coordinated X-ray/opt. obs.: [128]; UV obs.: [42, 163, 275, 594]; wind structure: [816]; mwl obs.: [731]; optical sp. type: [see 568]; distance of 3.1 kpc [552] and 1.4 kpc [322].						
1E 1145.1−6141	11 47 28.6	V830 Cen	13.1, 1,5, 0.15	9.607	4–70	14.4
	−61 57 14		B2Iae, 1.6	9.110		298
P	295.5, −0.01	[322]	[166, 331]	8.810	[322, 50]	[628, 414]
Pos. (o 2"): [322]; X-ray outburst: [50]; X-ray pulsations: [828]; period of 10.76 d from opt. sp.: [325]; system velocity: [795]; 15' away from 1145-619: [414, 829]; orbital elements (e = 0.2 and $f(m)$ = 5.1 $M_\odot$): [628]; distance of 8 kpc: [331].						
4U 1223−624	12 26 37.6	BP Cru ?	10.8, 1.76, 0.42	6.717	9–1000	41.59
GX 301−2	−62 46 13	WRA 977	B1–1.5 Ia, 1.8.4	6.077		696
P, T, C	300.1, −0.04	[65]	[57, 274, 764, 805]	5.672	[64]	[547, 689]
Pos. (o 3"): [64]; X-ray spectrum: [830]; recurrent outbursts: [615, 677, 820]; X-ray obs.: [262, 419, 423, 658, 826]; pulse-phase spectr.: [421]; pulse profile: [524]; X-ray dips: [422]; aperiodic X-ray variability: [35, 753]; rapid spin-up episodes: [393]; optical spectrum: [274, 323, 586, 764]; optical photom.: [796]; system velocity: [795]; IR obs.: [238]; a hypergiant (?): [368]; distance of 1.8 and 5.3 kpc: [368, 586]; variable cyclotron line (B of 3.4–5.8×10^{12}G): [406, 413, 570]; eccent. 0.46: [393, 689].						
IGR J12349−6434	12 34 54.7	RT Cru	12.46, 1.75, 0.28	6.646	1.3	
	−64 33 55.6		O–A,	5.583	(0.5–10 keV)	
T, P	301.2, −1.8		[96]	5.185	[777]	
Pos. (x 6"): [777]; discovery: [92]; optical ctpt. RT Cru?: [487]; accreting sys.: [487]; irregular variable: [96]; large-amplitude ($\triangle V = 3.8^m$) outburst and short-term flickering: [96]; symbiotic star: [96]; tentatively a HMXB.						
1H 1238−599	12 42 01.0				3–16	
3A 1239−599	−60 12 00					191
P	301.8, +2.7	[175]			[64, 815, 850]	[309]
Pos. (x 30"): [175]; hard X-ray spectrum: [309]; see also: [78].						
1A 1244−604	12 47 35				<24–100	
	−60 38 34					
T	302.5, +2.2				[78]	
Pos. (x 6.2'): [78].						
1A 1246−588	12 49 36				<24–300	
4U 1246−58	−59 07 18					
T	302.7, +3.7				[78]	
Pos. (x 4.5'): [78].						
1H 1249−637	12 42 50.3	HD 110432	5.31, 0.27, −0.79	4.655	2.2	
	−63 03 31	SAO 252002	B0.5IIIe, 0.40	4.339		14200?
	302.0, −0.2	BZ Cru [719]	[111, 564]	4.038	[850]	[771]
Pos. (o): [778]; Be star: [810]; UV obs.: [111]; interstellar absorption lines: [149]; optical obs.: [155, 212]; variable radial velocity: [73]; distance of 0.3 kpc: [94]; 14 ks period: [771]; a Be/WD candidate?: [771].						
1H 1253−761	12 39 14.6	HD 109857	6.49, 0.08, −0.24	6.169	0.6	
	−75 22 14	SAO 256967	B7 Vne, 0.20	6.169		
	302.1, −12.5	[719]	[111, 564]	6.102	[850]	
Pos. (o): [778]; Be star: [810]; visual double (sep. 2.2"): [202]; no optical brightness variations: [20]; white-dwarf accretor(?): [818]; distance of 0.24 kpc: [94].						

Table 1. (continued)

Name(s) type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
1H 1255−567	12 54 36.9 −57 10 07 303.4, +5.7	μ^2 Cru SAO 240367	5.17, −0.12, −0.51 B5 Ve, 0.04 [564, 710]	5.316 5.383 5.313	0.8 [850]	
Pos. (o): [778]; Be star: [810]; $v_r sini \sim 220 km/s$: [710]; visual double with μ^1 Cru: [299]; slow & small optical variability: [597]; distance of 0.11 kpc : [94].						
4U 1258−61 GX 304-1 P, T?	13 01 17.1 −61 36 07 304.1, +1.2	V850 Cen *2 (MMV) [489]	13.5–14.2, 1.7, 0.8 B0.7Ve, 2.0 [136, 568, 585]	9.798 9.297 9.040	0.3–200 [64, 601]	133? 272 [547, 617]
Pos. (IR): [489]; long-term X rays: [614, 617]; X-ray off state: [601]; X-ray spectrum: [658, 830]; opt. spectrum: [489, 764]; sp.-type B2V: [552]; long-term opt. variations: [136]; $v_r sini \sim 600$ km s^{-1}: [585]; system velocity: [795]; IR obs.: [238]; distance of 2.4 kpc: [585].						
2RXP J130159.6−635806 T, P	13 01 58.7 −63 58 09 304.1, −1.1		 [93]	13.0 12.1 11.4	6.3 (20–60 keV) [93]	 704 [93]
Pos. (IR): [93]; quick spin-up: [93]; distance 4–7 kpc: [93]; hard X-ray spectrum: [93]; a high mass X-ray binary with Be companion: [93].						
SAX J1324.4−6200 P	13 24 26.3 −62 00 53 306.8, +0.61		 	 	0.4 [9]	 170.84 [9]
Pos. (x 1.5'): [9]; distance >3.4 kpc: [9]; hard X-ray sp.: [9]; likely a Be/X-ray system: [9].						
2S 1417−624 P, T	14 21 12.9 −62 41 54 313.0, −1.6	*7 [13]	17.2:, 0.7:1.7, 0.8 B1Ve, 2: [258]	13.297 12.747 12.294	2–43 [64]	42.12 17.6 [207, 373]
Pos. (o): [64]; X-ray obs.: [13, 207, 373]; opt. id: [258]; orbital parameters (e= 0.45): [207]; Centaurus 1971–2 transient (?): [123]; 1994 II-type outburst: [207]; X-ray outburst: [333]; distance of 1.4–11.1 kpc: [258].						
SAX J1452.8−5949 P	14 52 49.3 −59 49 18 317.6, −0.44		 	 	0.04 [569]	 437.4 [569]
Pos. (x 50"): [569]; X-ray sp.: [569]; distance 6–12 kpc: [569]; probably a Be/X-ray binary: [569].						
4U 1538−52 P, E, C	15 42 23.3 −52 23 10 327.4, +2.2	QV Nor *12 [12]	14.4, 1.9, 0.6 B0 Iab, 2.1 [147, 330, 584]	10.358 9.910 9.677	<3–30 [64]	3.73 529 [31, 158]
Pos. (o): [64]; distance of 4.5 to 6.4 kpc: [100, 648]; X-ray obs.: [122, 459, 661]; X-ray eclipses: [31, 158]; X-ray orbit: [459]; X-ray spectrum: [459, 830]; X-ray cyclotron line at ~21 keV : [98, 662]; long-term spin-up trend: [679, 662]; eccentricity of 0.174: [99]; orbital decay -2.9×10^{-6} yr^{-1}: [99]; opt. light curve: [575]; radial-velocity curve: [147, 648]; $v_r sini \sim 200 km/s$: [147]; system velocity: [795]; opt. obs.: [584].						
XTE J1543−568 T, P	15 44 01 −56 45.9 325.0, −1.5		 	 	8 [475]	75.56 27.12 [342, 475]
Pos. (x 2'): [342]; a BeXRB at $d \geq 10$ kpc: [342]; orbital parameters (e<0.03 and f(M)=8.2$M_\odot$): [342].						
2S 1553−542 P	15 57 49.0 −54 24 54 327.9, −0.9		 	 	27 [64]	30.6 9.3 [375, 547]
Pos. (x 3.5"): [64]; opt. counterpart likely a Be star: [375]; X-ray orbit: [375]; e<0.09: [375].						

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
1H 1555−552	15 54 21.8 −55 19 45 327.0, −1.2	HD 141926 SAO 243098 [719]	8.60, 0.56, −0.43 B2IIIn, [192, 806]	7.098 6.841 6.509	1.7 [850]	
Pos. (o): [778]; Be star: [810]; distance of 0.96 kpc: [257]; Herbig Ae/Be star?: [806]; low luminosity BeXRB: [771].						
IGR J16195−4945 AX J1619.4−4945	16 19 29 −49 45.5 333.5, 0.3	HD 146628? [707]	10.08, 0.44, B1–2Ia, [707]	8.575 8.412 8.377	9 [707]	
Pos. (x 1'): [707]; optical ctpt HD 146628?: [707]; variable, highly absorbed hard spectrum: [707]; likely a neutron star in a HMXB: [707]; distance of 7 kpc: [707].						
IGR J16283−4838	16 28 10.7 −48 38 55 335.3, 0.10	 [33]		16.798 15.815 13.948	13 (20–40 keV) [33]	
Pos. (x 5"): [33]; highly absorbed (variable) and flat X-ray spectrum: [33]; possible IR counterpart in the 2MASS and Spitzer survey: [33]; a HMXB containing a NS: [33].						
IGR J16318−4848	16 31 48.31 −48 49 00.7 335.6, −0.4	* 1 [205]	>21.1, , , 5.6 [205]	10.33 <10.4 <9.1	45 [498]	
Pos. (o): [205]; the first source discovered by INTEGRAL: [143, 813]; very high absorption ($N_H > 10^{24}$ cm^{-2}): [813, 205, 498]; very strong Fe Kα emission line and significant Kβ and Ni Kα lines: [498]; IR ctpt.: [216]; previous ASCA outburst in 1994: [543]; I=16.05, R=17.72: [205]; a HMXB with a B0–B5 sgB[e] at a distance between 0.9–6.2 kpc: [205].						
IGR J16320−4751 AX J1631.9−4752 P	16 32 01.9 −47 52 27.1 336.3, 0.17	* 1 [671]		14.080 13.028 10.986	17 [343]	8.96 1309 [142, 450]
Pos. (x 4"): [671]; discovery: [767]; heavily absorbed: [669, 343]; persistent for at least 8 years: [343]; star '1' IR ctpt: [671]; SG/X-ray binary?: [343].						
IGR J16358−4726 T, P	16 35 58 −47 24.4 337.1, −0.00				7 [591]	 5880 [591]
Pos. (x 2'): [44]; no radio emission detected; heavily absorbed source: [44]; probably a HMXB with spin 5880 s: [591]; Fe Kα emission line: [591]; rough distance of ~6–8 kpc: [645].						
AX J1639.0−4642 IGR J16393−4643 P	16 39 05.4 −46 42 14 338.01, 0.07			14.633 13.316 12,780	0.8 (0.7–10keV) [51]	 912.0 [51]
Pos. (IR): [51]; IR ctpt 2MASS J16390535−4642137; huge N_H, hard spectral index, probably a HMXB: [742, 120, 51]; well within the error of 3EG J1639−4702: [120]; large variations in intensity: [51]; X-ray lines: [51]; the radio ctpt [120] out of the XMM error: [51].						
IGR J16418−4532	16 41 45 −45 31 12 339.2, 0.5				1 (20–40 keV) [446]	
Pos. (x 2'): [768]; in the Norma Region: [768]; heavily absorbed $N_H \sim 10^{23} cm^{-2}$: [410]; a HMXB?: [410].						

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
IGR J16465−4507	16 46 35.26			10.537	16	
	−45 07 04.5		B0.5I,	10.079	(18–60 keV)	228
T, P	340.1, 0.13		[560]	9.843	[446]	[447]
Pos. (IR): [560]; B = 15.2, R = 13.0: [714]; fast X-ray transient: [446]; extremely absorbed ($N_H \sim 7\times10^{23} cm^{-2}$): [447]; very high rotational velocity: [560]; 2MASS source (J=10.54) and No. 0448-0520455 in the USNO B1.0 catalog; distance of 12.5 kpc: [714].						
IGR J16479−4514	16 48 07.0		∼20.4, ,	12.954	470	
	−45 12 05.8			10.825	(20–30 keV)	
T	304.2, −0.12		[379]	9.798	[704]	
Pos. (x 6”): [379]; fast transient: [704]; a faint source with V∼20.4: [379]; $N_H = 1.2\times10^{23} cm^{-2}$: [447]; IR source IRAS 16441−4506 in the error: [704]; outburst or type I burst?: [704]; a NS in a HMXB?: [447, 704].						
IGR J16493−4348	16 49 21				0.4	
	−43 48 36					
	341.4, 0.6				[472]	
Pos. (x 2’): [253]; in the Norma Region: [253]; only 45” off the radio pulsar PSR J1649−4349: [253]; variable on time scales of 100 to 1000 s: [472]; no significant pulsations: [472]; heavily absorbed power law ($N_H \sim 10^{23} cm^{-2}$ and α=1.4): [472]; a HMXB?: [410].						
OAO 1657−415	17 00 48.9	star		14.665	4–42	10.4
	−41 39 21.6		B SG, 6.6	13.778		38
P, E	344.4, 0.31	[83]	[83]	13.383	[64]	[81, 547]
Pos. (x 0.5”): [83]; V861 Sco not the optical ctpt.: [14]; X-ray obs.: [366]; X-ray spectrum: [830]; hard X-ray obs.: [509]; X-ray eclipsing: [81]; orbital solution: [81]; highly reddened: [83]; distance of 6.4 kpc: [83]; iron K emission line: [83].						
4U 1700−37	17 03 56.8	HD 153919	6.6, 0.27, −0.72	5.744	<11–110	3.41
	−37 50 39	V884 Sco	$O6.5Iaf^+$, 0.52	5.639		
E, C	347.8, +2.2	[362]	[276, 596, 848]	5.496	[64]	[362]
Pos. (o): [64]; eclipsing system: [362]; X ray during strong flares and eclipse: [789]; X-ray variab.: [35, 174, 265]; hard X-ray sp.: [600]; no X-ray pulsations: [174, 246]; X-ray sp. lines: [789, 58]; QPO?: [58]; broad cyclotron absorption feature at ∼37 keV?: [649]; opt. light curve: [19, 799]; opt. sp.: [125, 154, 191, 287]; radial-velocity curve: [174, 273, 277]; e = 0.22: [277]; $v_r sini \sim$ 140–300 km s^{-1}: [125, 154, 313, 848]; system velocity: [795]; UV obs.: [181, 276, 367]; soft X-ray Raman scattering: [367]; opt. polarimetry: [171, 172]; polarization: [849]; IR obs.: [238, 596]; system parameters: [281]; possibly 13.81 days period: [395, 303]; runaway from OB association SCO OB1: [10]; decrease in orbital period: [265]; physical parameters with $\dot{M} = 9.5\times10^{-6}$ $M_\odot$/yr, $M_* = 58M_\odot$and $M_x = 2.44M_\odot$: [106]; low mass BHC?: [649]; distance of 1.9 kpc: [10].						
AX J1700−419	17 00 5.3				0.4	
AX J170006−4157	−41 57 44					714.5
P	344.0, +0.25				[770]	[770]
Pos. (x 1’): [770]; a WD binary also possible: [770].						
IGR J17091−3624	17 09 02.3				22	
	−36 23 33				(15–40 keV)	
T, R	349.5, 2.2				[411]	
Pos. (r): [680]; radio ctpt: [680]; radio obs., microquasar?: [577]; X-ray large variability: [448, 411]; BHC: [448]; uncertain if a Be/X-ray binary or LMXB: [343].						
EXO 1722−363	17 25 11.4			14.218	0.2–8	9.74
IGR J17252−3616	−36 16 57.5			11.811		413.9
P, E	351.5, −0.35			10.672	[754, 141]	[141, 754]
Pos. (IR): [861]; 2MASS J17251139-3616575: [861]; apparent total eclipses: [141]; hard X-ray spectrum: [754]; very large N_H: [749, 861]; Fe Kα line at 6.4 keV: [861]; at Galactic bulge region: [749]; mass function: [141]; a SG/X-ray binary: [141, 861].						

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
XTE J1739−302 IGR J17391−3021 T	17 39 11.58 −30 20 37.6 358.1, 0.45		14.89, 2.97, 1.68 O8Iab(f), 3.4 [562]	8.600 7.823 7.428	136 (2–25 keV) [715]	 1500? [684]
Pos. (x): [718]; fast transient: [715, 718]; USNO A2.0 (0525 28760590) and 2MASS source: [718]; highly reddened: [562]; radio obs.: [295]; radio source 2' away from the Chandra source: [681]; X-ray spec: [715]; distance of ∼2.3 kpc: [562].						
RX J1739.4−2942 GRS 1736−297 T	17 39 30.1 −29 42 07 358.6, −0.7				2.5 (3–12 keV) [593]	
Pos. (x 19.2"): [540]; hard X-ray spec.: [540]; X-ray observation: [593, 842]; a Be/X-ray binary instead of LMXB?: [540]; LMXB in SIMBAD.						
AX J1740.1−2847 P	17 40 11.6 −28 47 48 359.5, 1.1				0.2 [683]	 729 [683]
Pos. (x 30"): [683]; possibly a HMXB near the Galactic center region: [683].						
RX J1744.7−2713	17 44 45.7 −27 13 44 1.36, +1.1	V3892 Sgr HD 161103	8.4, 0.44, −0.64 B0.5V–IIIe, [180, 727]	7.054 6.843 6.507	0.06 [539]	
Pos. (o): [727]; strong emission lines: [539]; distance 0.8 kpc: [539]; IR photom. and spectrum: [724, 308]; *vsini* = 224 km s^{-1}: [727].						
AX J1749.1−2733 T	17 49 06 −27 33 1.58, 0.06				0.2 (0.7–10 keV) [684]	
Pos. (x ?): [684]; fast X-ray transient: [684, see also 334]; HMXB?: [560].						
AX J1749.2−2725 T, P	17 49 10.1 −27 25 16 1.7, +0.1				1.36 [769]	 220.38 [769]
Pos. (x 1'): [769]; weak X-ray source: [769]; hard X-ray sp.: [769, 684]; likely a BeXRB with NS: [769]; within error of a nearby WD: [769]; distance 8.5 kpc: [769].						
GRO J1750−27 P, T	17 49 12.7 −26 38 36 2.4, +0.5				56 [698]	29.8 4.45 [698]
Pos. (x 4"): [698]; no optical counterpart reported: [698]; likely Be system: [698]; large spin-up: [698]; orbital parameters (e=0.36): [698]; distant source (18 kpc?): [698].						
IGR J17544−2619 1RXS J175428.3−262035 T	17 54 25.28 −26 19 52.6 3.2, −0.34	* C1 [595]	12.65, 1.79, O9Ib, 2.07 [595]	8.791 8.310 8.018	89 (17–45keV) [250]	
Pos. (IR 0.2"): [668]; short X-ray flare: [743, 250]; 2MASS J17542527-2619526 and also in the USNO B1.0 Catalogue: [668]; ultra-soft quiescent X-ray: [334]; a NS: [334]; distance of 8 kpc or between 3–4 kpc: [243, 595].						
SAX J1802.7−2017 IGR J18027−2016 T, P, E	18 02 39.9 −20 17 13.5 9.4, 1.04				1.5 (0.1–10 keV) [18]	4.6 139.61 [18]
Pos. (x 2'): [18]; ∼ 22′ away from the bright X-ray source GX 9+1: [18]; system parameters ($e \leq 0.2$ and $f(M) \sim 17 \pm 5 M_{\odot}$): [18, 291]; an eclipsing HMXB (at distance of 10 kpc?): [18, 291].						

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
4U 1807−10	18 11 −10 54				<2–10	
T	18.6, +3.9				[215]	
Pos. (x 1.3°): (large error box): [215]; uncertain if HMXB or LMXB.						
SAX J1818.6−1703	18 18 39 −17 03 06	HD 168078?	10.7, −0.1, B3III,		56 (2–9 keV)	
T	14.1, −0.73		[338]		[338]	
Pos. (x 3'): [338]; fast transient: [704]; optical ctpt HD 168078?: [338].						
SAX J1819.3−2525 XTE J1819−254	18 19 21.48 −25 25 36.0	V4641 Sgr	8.8–13.5, −0.2, 1.8 B9III, 0.32	12.532 12.364	1–13000 (2–12 keV)	2.8
T, R	6.8, −4.8	[244]	[572, 741]	12.27	[713]	[572]
Pos. (o 0.7"): [686]; misidentified as GM Sgr: [686]; X-ray light curve: [339]; strong relativistic Fe-Kα emission line: [339, 515]; X-ray flare sp.: [339, 457]; fast and large X-ray outbursts: [713, 834]; X-ray sp. harder during decay and considerably softer in quiescent state: [834]; opt. outburst: [741, 87]; opt. sp.: [572]; radio obs., highly relativistic jet (μQ): [298]; orbital period: [370, 572]; mass of the compact star 9.6 $M_\odot$: [572]; distance between 7–12 kpc: [87, 572]; super-Eddington outburst: [646].						
AX J1820.5−1434	18 20 29.5 −14 34 24		O9.5–B0Ve		8	152.26
P	16.5, +0.07		[350]		[449]	[382]
Pos. (x 0.5'): [382]; X rays: [449]; R = 17.33: [350]; a highly obscured accreting binary: [382]; distance of 4.7 kpc: [382].						
RX J1826.2−1450	18 26 15.06 −14 50 54.3	LS 5039	11.23, 0.94, −0.16 ON6.5V((f)), 1.28	9.024 8.751	0.3	3.90
R	16.9, −1.3	[539]	[504, 539]	8.604	[539]	[80]
Pos. (r): [653]; distance 2.5 kpc: [80]; small optical photom. var.: [482]; radio ctpt.: [479]; X-ray and radio obs.: [651, 652]; X-ray variability: [60, 638]; relativistic jet, γ-ray-emitting persistent (μQ): [582, 583]; orbital X-Ray variability: [60]; low X-ray state: [484]; optical polarimetry: [121]; orbital parameters: [504, 80]; e = 0.35 and M_X = 3.7 (a BHC?): [80]; runaway: [653, 503]; very high energy γ rays: [5].						
XTE J1829−098	18 29 44.0 −09 51 23				4	7.82
T, P	21.7, 0.28				[471]	[471]
Pos. (x 4"): [268]; I = 14.4: [269]; heavily absorbed ($N_H \sim 10^{23} cm^{-2}$): [471]; hard spectrum favoring a HMXB: [471]; 8" away from a very bright SG star in 2MASS (with J=8.04, H=6.11, K=4.98): [269].						
H 1833−076 Sct X-1	18 36 28.6 −07 36 21				1.6–200	111
P, T	24.5, −0.16	[247]			[399, 618, 815]	[403, 547]
Pos. (x 30"): [630]; alternative position: [630]; X-ray obs.: [86, 292]; X-ray pulsations: [403]; hard X-ray spectrum: [129, 630]; long-term X-ray record: [618]; transient: [129].						
AX J1838.0−0655	18 38 01 −06 55 54 25.2, −0.19				0.5 (0.7–10keV) [742]	
Pos. (x 1.0'): [742]; $N_H = 1.1 \times 10^{23} cm^{-2}$, probably a HMXB: [742]; a SNR or a pulsar with nebula: [462]; high-energy cutoff value above 30 keV (BHC?): [447].						
GS 1839−04	18 41 48 −04 27				2.5	81.1
P, T	27.9, +0.11				[399]	[399]
Pos. (x 24'): [399].						

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
GS 1839−06	18 41 42				1	
	−05 51					
T	26.6, −0.5				[399]	
Pos. (x 30'): [399].						
AX J1841.0−0536	18 41 00.43			9.74	4	
IGR J18410−0535	−05 35 46.5			9.22		4.7394
T, P	26.8, −0.24			8.93	[22]	[22]
Pos. (o): [271]; fast transient: [22]; B = 15.91, R = 12.78: [271]; distance 10 kpc?: [22, 271]; 6.4 keV line: [22]; heavily absorbed hard spectrum: [22, 271]; weak Hα emission: [271].						
GS 1843+009	18 45 36.8	star	20.2,		33	
	00 51 48.3		B0–B2IV–Ve, 2.3–2.9			29.5
P, T, C	33.0, 1.7	[351]	[351]		[399]	[399]
Pos. (o): [351]; X-ray obs.: [399, 400, 603]; hard X-ray outburst: [845, 89]; possible cyclotron feature (B = 1.7×10^{12}G): [512]; V−R = 2.1: [351]; distance ≥10 kpc: [351]; strong Hα emission: [351].						
2S 1845−024	18 48 18				1–44	241
GS 1843−02	−02 25 12					94.8
P, T	30.4, −0.4	[247]			[64, 399]	[859, 399]
Pos. (x 30"): [64]; also known as 4U 1850-03 and GRO J1849-03; X-ray obs.: [177, 399, 401, 700]; distance: of 10 kpc: [399]; likely a Be/X-ray: [720, 209]; X-ray outburst: [209]; high eccentricity (0.88): [209].						
AX 1845.0−0433	18 45 02		13.96, 2.2, 1.04		45	
IGR J18450−0435	−04 33 31		O9.5I, 2.45		(0.7–10 keV)	
T	28.8, −1.22	[119]	[119]		[854]	
Pos. (x 1'): [119]; fast transient: [854]; discovery of x-ray source: [854]; optical spectrum: [119]; distance 3.6 kpc: [119].						
IGR J18483−0311	18 48.3				22	
	−03 11				(15–40 keV)	
T	29.8, −0.7				[91]	
Pos. (x 2'): [91]; in the Scutum arm, probably a HMXB: [343].						
GS 1855−02	18 58				2	
	−02 44					
T	31.2, −2.7	[132]			[399]	
Pos. (x 10'): [399].						
XTE J1855−026	18 55 31.3				6	6.067
	−02 36 24					361
P, T, E	31.1, −2.1	[132]			[139]	[139]
Pos. (x 12"): [132]; iron emission line: [139]; most likely a SG/X-ray system: [139, 132]; BD-2 4786 not the opt. cpt.: [139]; X-ray sp.: [139]; within the error of src No. 6 in [399]: [139]; an extended near-total eclipse system: [132]; orbital parameter (e<0.04): [132]; consistent with a B0I star: [132]; highly absorbed: [132]; an early-type star (B=16.1, I=11.5) right on the edge of the ASCA error: [803]; distance ∼ 10 kpc: [139].						
XTE J1858+034	18 58 36	star?	18.0, 1.6,		25	
	03 26 09					221.0
P, T	36.8, −0.02	[640]	[640]		[643]	[748]
Pos. (o): [640]; 0.11 Hz QPO: [592]; outburst: [529]; very hard X-ray sp.: [592, 643]; probably a Be/X-ray sys.: [748]; emission-line optical ctpt. though out of the JEM-X error: [640].						

Table 1. (continued)

Name(s) Type	RA (J2000) DEC (J2000) l^{II}, b^{II}	Opt. Ctp. [FC]	V, $B-V$, $U-B$ Sp. type, E_{B-V}	J H K	F_x μJy	P_{orb}(d) P_{pulse}(s)
XTE J1859+083	18 59.1				8	
	08 15					9.801
T, P	41.1, 2.1				[474]	[474]
Pos. (x 2'): [474]; X-ray pulsar: [474]; tentatively a HMXB.						
XTE J1901+014	19 01 39.90	star	18.72, 2.44,		500	
1RXS J190141.0+012618	01 26 39.2				(2–12 keV)	
T	35.4, −1.6	[612]	[612]		[644]	
Pos. (o): [612]; fast transient: [644].						
4U 1901+03	19 03 37.1				<2–333	22.58
	+03 11 31					2.763
P, T	37.2, −1.25	[224]			[123, 224]	[224]
Pos. (x 1'): [224]; transient 1971: [123, 214]; hard X-ray spectrum: [123]; distance of 10 kpc?: [224]; the longest (32.2 yr) recurrence X-ray transient: [224]; orbital parameters (e = 0.036 and F_x(M)=2.58$M_\odot$): [224]; lack of bright IR ctpt.: [224]; unlikely SG companion: [224]; accreting from the wind of OB main-sequence star?: [224].						
XTE J1906+09	19 04 47.48	star		15.18	0.3–5	
	09 02 41.8			14.17		89.17
P, T	42.5, 1.2	[240]		13.50	[839]	[473]
Pos. (o): [240]; R=18.7: [240]; distance ≥ 10 kpc: [473]; likely a Be/X-ray binary: [473, 839, 240]; X-ray outbursts: [473, 839, 240]; orbital period between 26–30 d: [839]; low eccentricity (0.03–0.06): [839].						
4U 1907+09	19 09 39.3	star	16.4, 3.2,		4–275	8.38
H 1907+097	+09 49 45		O8–9 Ia, 3.45			438
P, T, C	43.7, +0.5	[696]	[144, 696]		[64]	[458, 477]
Pos. (o): [696]; distance ∼5 kpc: [144]; long-term X-ray record: [618]; 1980 outburst: [477]; cyclotron lines at ∼19 and ∼39 keV (B of 2.1×10^{12} G); [150]; heavily reddened massive star with broad Hα emission: [696]; orbital parameters: [127, 458]; orbital modulation X-ray flux: [477]; X-ray dip due to cessations of the mass accretion rather than occultation: [336]; long-term spin-down: [337]; e = 0.28: [337]; opt. spectrum: [353, 791]; $v_r sini$ ∼ 85 km s^{-1}: [791].						
3A 1909+048	19 11 49.6	SS 433	14.2, 2.1, 0.6	9.398	2–10	13.1
	+04 58 58	V1343 Aql	pec, 2.6	8.739		
E, R	39.7, −2.2	[425]	[466, 546, 811]	8.163	[64]	[146]
Pos. (o,r): [64]; X-ray obs.: [23, 35, 68, 69, 259, 372]; Doppler shift X-ray Fe line: [497, 822]; extended X-ray lobes: [701, 821]; γ-ray obs.: [226, 415]; Doppler shifted opt. emission lines: [464, 467]; high-speed jet velocity (v∼0.26) (μQ): [4, 235]; stationary emission lines from accretion disk: [204]; kinematic model with precessing of 164 d: [4, 183]; opt. spe.: [190, 193, 397, 468]; opt. spectrophot.: [6, 396, 811]; opt. lt. curve: [16, 288, 424, 426, 427]; long-term optical phot.: [288, 377, 499]; X-ray/opt. eclipse: [732]; IR obs.: [390, 814]; precession clock: [7]; radio structure: [24, 195, 196, 673, 722, 801]; radio variability: [56, 203]; distance 4.61 in [736], 5.5 kpc in [296, 49]; nature of the compact star: [170]; circularly polarized radio emission: [199]; BH with a mid-A SG sys.: [293, 90]; evolutionary status: [381]; absorption features from accretion disc wind?: [25]; sharp variations in the X-ray, IR and radio, double X-ray Fe line profiles from the jet: [85]; massive ejection event with a QPO-like feature near 0.1 Hz: [398]; extended radio emission: [48]; X-ray dominated by precessing jets: [551, 476]; bullet-like jets: [84]; highly ionized Doppler-shifted iron emission lines: [511]; identifiable deviations from the traditional kinematic model: [49]; rapid chaotic optical variability: [647].						
4U 1909+07	19 10 48.2	* A		13.228	1–7	4.4
X1908+075	+07 35 52.3		OI7.5–9.5	11.457		604.68
P	41.9, −0.81	[533]	[533]	10.480	[824]	[824, 428]
Pos. (IR): [533]; distance 7 kpc: [533]; highly-absorbed: [824]; orbital inclination angle (38°–72°): [428]; e=0.021, f(M)=6.07$M_\odot$: [428].						

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
IGR J19140+0951	19 14 04.2			8.55	75–100	13.558
IGR J19140+098	9 52 58.3			7.67	(1–20 keV)	
	44.3, −0.46	[344]	[344]	7.06	[670]	[140]

Pos. (x 0.6"): [344]; also known as EXO 1912+097: [443]; a HMXB with a NS: [670]; variable absorption column density: [670]; bright iron line: [670]; IR ctpt. 2MASS 19140422+0952577: [344].

1H 1936+541	19 32 52.3	DM +53°2262	9.8, ,	10.231	0.7	
	+53 52 45		Be,	10.166		
	85.9, +15.9		[778]	9.975	[850]	

Pos. (o): [778]; Be star: [810]; X-ray emission not detected: [771]; consistent with a quiescent Be/NS sys.: [771].

XTE J1946+274	19 45 39.3	*A	16.9, 1.72,	12.539	< 1–25	169.2
GRO J1944+26	27 21 55.4		B0–1 IV-Ve, 1.9–2.2	11.831	(2–60 keV)	15.8
P, T, C	63.2, +1.4	[802]	[802]	11.333	[745]	[841, 711]

Pos. (o): [802]; transient Nov-Dec 1976: [815]; X-ray flare: [76]; ~80 day period: [76]; very hard X-ray spec.: [711]; distance of 9.5 kpc: [802, 841]; $e = 0.33$ and $f(M) = 9.7M_{\odot}$: [841]; IR excess and strong Hα emission: [802]; cyclotron line at 36.2 keV ($B = 4.2 \times 10^{12}$ G): [283].

KS 1947+300	19 49 30.5	*3	14.24, 0.93, −0.09	11.67	< 10–375	40.4
GRO J1948+32	+30 12 24		B0Ve, 1.09	11.43		18.7
P, T	66.1, +2.1	[248]	[559]	11.21	[59]	[82, 223]

Pos. (o): [248]; Hα emission: [245, 559]; IR photom. (J=11.67, H=11.43, K=11.21): [559]; distance about 10 kpc: [559]; eccentricity 0.033: [223]; mass function: [82]; frequency glitch: [223].

4U 1956+35	19 58 21.7	HD 226868	8.9, 0.84, −0.26	6.872	235–1320	5.
Cyg X-1	+35 12 06	V1357 Cyg	O9.7 Iab, 1.06	6.652		
U, R	71.3, +3.1	[624]	[124, 157, 564]	6.501	[64]	[231]

Pos. (o,r): [64]; hard X- and γ- ray obs.: [230, 432, 685, 697, 779, 780]; long-term X-ray obs.: [301, 620, 239, 610, 857, 835]; X-ray low/high states: [431, 751]; X-ray dips: [21, 384, 642]; rapid X-ray variability: [34, 35, 442, 505, 525, 526, 527, 678]; QPO: [219, 808]; chaotic (?): [441, 785]; 300 d period: [232, 376, 378, 620]; X-ray spectrum: [26, 27, 188, 385, 445, 809]; γ-ray obs.: [433, 599]; radio obs.: [66, 67, 294, 297]; opt. spectra: [1, 124, 233, 234, 565]; radial-velocity curve: [231, 232, 565]; $v_r sini \sim 100$ km s^{-1}: [233]; system velocity: [795]; nature of the compact star: [52, 705, 823]; opt. light curve: [378, 453]; opt. UV polarimetry: [173, 218, 849]; UV obs.: [157, 181, 774, 852]; IR obs.: [420]; fund. parameters, $M_* = 17.5M_{\odot}$and $M_x = 10.1M_{\odot}$: [290]; massive BH formed without SN: [521]; dark jet: [222]; precession of the inner accretion disk: [773]; model constraint on the masses: [860]; focused companion wind: [517]; quenched millimeter emission: [765]; short cutoff timescales: [201]; distance about 2.14 kpc: [491]; broad-band X-ray spectra: [329]; giant outburst: [242]; powerful millisecond flares: [229]; radio jets with a speed in excess of 0.6c (μQ): [735]; orbital elements: [236]; probing the inner region: [168]; X-ray/optical sp.: [752]; composite Fe Kα emission line: [516]; short-time-scale correlations between line and continuum fluxes: [455]; soft Gamma-ray spectral variability: [501]; high-resolution X-ray spectrum: [695]; temporal evolution of X-ray lags: [609]; very flat radio-millimetre spectrum: [200]; X-ray sp.: [237]; orbital modulation: [607].

W63 X-1	20 19				1.5	
2E 2017.1+4553 ?	+45 42				(0.3–4.5 keV)	36
P	82.3, 5.4				[703]	[650]

Pos. (x ?): [846]; in the northern part of SNR W63 [846] (G82.2+5.3), possibly associated with the SNR: [650]; distance between 1.3–1.9 kpc: [674]; much harder than diffuse soft X-ray emission from SNR W63: [650]; possible counterpart having Hα excess emission: [650]; HMXB with a Be companion: [650].

EXO 2030+375	20 32 15.2	*2	19.7, 3.3,	12.05	<0.5–1400	46.02
	+37 38 15	V2246 Cyg	B0Ve, 3.8	10.835		41.8
P, T	77.2, −1.3	[114, 535]	[114, 535]	10.074	[587]	[547, 587, 739]

Pos. (o 2"): [535]; giant outburst: [587]; rapid X-ray variability: [35]; 0.2 Hz QPO: [8]; opt. spe.: [114, 357]; long-term opt./IR variability: [635]; distance of 7.1 kpc: [840]; eccentricity of 0.41: [840]; X-ray outburst at apastron: [417]; transition from global spin-down to spin-up: [843].

Table 1. (continued)

Name(s)	RA (J2000)	Opt. Ctp.	V, $B-V$, $U-B$	J	F_x	P_{orb}(d)
	DEC (J2000)		Sp. type, E_{B-V}	H	μJy	P_{pulse}(s)
Type	l^{II}, b^{II}	[FC]		K		
4U 2030+40	20 32 26.5	V1521 Cyg		15.309	90–430	0.20
Cyg X-3	+40 57 09		WR star, 6.3	13.192		
R	79.8, +0.7	[812]	[792]	11.921	[64]	[590]
Pos. (r): [64]; I=20.0: [812]; distance about 10 kpc: [167]; orb. period change: [386, 787, 709]; long-term X-ray obs.: [301, 619]; no X-ray pulsations: [386, 851]; orbital X-ray curve: [54, 383]; aperiodic variability: [35, 61]; X-ray halo: [530]; transient QPO: [788]; radio outbursts: [255, 360, 567]; quiescent radio flares (period 4.95 h?): [531]; two-sided relativistic jet (μQ): [481, 480, 740]; one-sided highly relativistic jet: [225, 520]; IR phot.: [32, 488, 490]; IR spectra: [371, 792]; X-ray spectrum: [383, 825]; sp. variability during outburst and quiescence: [198]; VLBI obs. during outburst: [691]; intense stellar wind: [613, 790]; correlated behavior between hard X-ray and radio: [500]; hot spots excited by jets: [483]; hard X-ray time lag: [95]; multifrequency radio observations: [518]; extended X-ray emission: [284]; γ-ray obs.: [563]; Doppler shifts of the X-ray emission lines: [723]; high-resolution X-ray spectroscopy: [574].						
RX J2030.5+4751	20 30 30.8	SAO 49725	9.27, 0.38, −0.65	8.058	0.04	
	47 51 51	BD +47 3129	B0.5V–IIIe,	7.854		
	85.2, +5.0	[539]	[539]	7.624	[539]	
Pos. (o): [300]; strong emission lines: [539]; distance 2.2 kpc: [539]; likely Be/X-ray binary: [539].						
GRO J2058+42	20 58 47.5	star	14.92, 1.12,	11.74	320	55.03
CXOU J205847.5+414637	41 46 37		O9.5–B0IV–Ve, 1.4	11.282		198
P, T	83.6, 2.7	[844]	[640]	10.930	[837]	[837, 844]
Pos. (o): [640]; giant outburst in 1995: [837]; distance 9.0 kpc: [640]; Chandra obs.: [844]; optical ctpt to the Chandra src 3" outside the GRO error: [844]; strong Hα emission line: [844]; 2MASS source: [844]; pulsations not detected with Chandra: [844].						
SAX J2103.5+4545	21 03 35.7	star	14.2, 1.12,	11.842	20	12.68
	45 45 04		B0Ve, 1.36	11.535		358.61
P, T	87.12, −0.7	[639]	[639]	11.362	[30]	[30, 310]
Pos. (o): [639]; HD200709 not opt. cpt.: [30]; X-ray spe.: [310]; large spin-up: [708]; 0.044Hz QPO: [332]; eccentricity 0.4±0.2: [30]; distance of 6.5 kpc: [639]; fast optical spectral variability: [639]; *vsini* = 240km s^{-1}: [639].						
1H 2138+579	21 39 30.6	V490 Cep	14.29, ,	11.829	<6–100	
Cep X-4	+56 59 12.9		B1–B2Ve, 1.3	11.414		66.2
P, T, C	99.0, 3.3	[53]	[53, 53]	10.926	[402]	[402, 547]
Pos. (o): [667]; X-ray obs.: [402]; X-ray cyclotron line (B=2.5×10^{12} G): [513]; same as Cep X-4(?): [470, 782]; X-ray outburst: [838]; orbital period 23–147.3 days: [838]; distance of 3.8 kpc: [53]; *vsini* 460 km s^{-1}: [53].						
1H 2202+501	22 01 38.2	BD +49°3718	8.8, 0.17,	8.778	0.7	
	+50 10 05	V2175 Cyg	Be, 0.30	8.701		
	97.2, −4.0	[719]	[778]	8.496	[850]	
Pos. (o): [778]; Be star: [810]; distance of 0.7 kpc: [94].						
4U 2206+543	22 07 56.2	BD +53 2790	9.85, 0.2, −0.6	9.218	0.6–5.5	9.57
3A 2206+543	+54 31 06		O9.5Ve, 0.5	9.116		
C	100.6, −1.1	[728]	[728, 555]	9.038	[815, 850]	[134, 654]
Pos. (o): [300]; X rays: [688]; NS system: [772, 485, 46, 654]; Be/WD system: [134]; distance 2.6 kpc: [47]; 392 s pulsation? [688] not true: [134]; cyclotron resonant feature at 32 keV (B=3.6×10^{12} G): [772, 485, 46]; not typical Be, but peculiar O9.5V star with relatively strong wind: [46, 555]; very low wind terminal velocity of ∼350 km s^{-1}: [654]; moderate eccentricity around 0.15?: [654]; wind-fed HMXB with ms donor: [654].						
SAX J2239.3+6116	22 39 20.90	star	15.1, 1.4,	11.450	16	262
1H 2214+589	+61 16 26.8		B0V-B2IIIe, 1.4	10.955		1247
T, P	107.7, +2.3	[340]	[340]	10.557	[340]	[340, 341]
Pos. (o 0.3"): [340]; distance of 4.4 kpc: [340]; position coincided with 4U 2238+60 and 3A 2237+608: [340].						

References

Bird, A.J., Barlow, E.J., Bassani, L. et al. 2004, ApJ, 607, L33
Caraveo, P.A., Bignami, G.F., Goldwurm, A. 1989, ApJ, 338, 338
Chaty, S., Charles, P. A., Martí, J. et al. 2003, MNRAS, 343, 169
Clark, G.W., Schmidt, G.D., Angel, J.R.P. 1975, IAUC 2843
Coe, M.J. 2000, ASPC, 214, 656
Coe, M.J. Fabregat, J., Negueruela, I. et al. 1996, MNRAS, 281, 333
Gotthelf, E.V. & Vasisht, G. 1998, NewA, 3, 293
Grimm, H.-J., Gilfanov, M., Sunyaev, R. 2003, MNRAS, 339, 793
Halpern, J.P., Gotthelf, E.V., Leighly, K.M., Helfand, D.J. 2001, ApJ, 547, 323
Halpern, J.P., Gotthelf, E.V., Helfand, D.J. 2004, ATel 289
Hang, H.R., Liu, Q.Z. & Xia, J.P. 1999, A&A, 344, 614
in't Zand, J.J.M. 2005, A&A, 441, L1
in't Zand, J.J.M., Halpern, J., Eracleous, M. et al. 2000, A&A, 361, 85
Kaper, L., van der Meer, A., Tijani, A.H. 2004, RMxAC, 21, 128
Kuntz, K.D., Gruendl, R.A., Chu, Y.-H. et al. 2005, ApJ, 620, L31
Kuulkers, E. 2004, in Proc. The Restless High-Energy Universe, ed. E.P.J. van den Heuvel et al., Nucl. Phys. B, 132, 466
Kuulkers, E. 2005, astro-ph/0504625
Lewin, W.H.G. & van der Klis, M. (eds), 2006, Compact Stellar X-Ray Sources, Cambridge University Press
Liu, J.-F., Bregman, J.N., Seitzer, P. 2004, ApJ, 602, 249
Liu, Q.Z. & Mirabel, I.F. 2005, A&A, 429, 1125
Liu, Q.Z., van Paradijs, J. & van den Heuvel, E.P.J. 2000, A&AS, 147, 25 (3rd edition)
Liu, Q.Z., van Paradijs, J. & van den Heuvel, E.P.J. 2005, A&A, 442, 1135
Markwardt, C.B., Swank, J.H., Smith, E. 2005, ATel 465
Marshall, F.E., in't Zand, J.J.M., Strohmayer, T. et al. 1999, IAUC 7240
Masetti, N., Bassani, L., Bird, A. J., Bazzano, A. 2005, ATel 528
Masetti, N., Orlandini, M. Palazzi, E. et al. 2006, astro-ph/0603227
Mereghetti, S., Belloni, T., Shara, M., Drissen, L. 1994, ApJ, 424, 943
Mereghetti, S., Israel, G.L., Stella, L. 1998, MNRAS, 296, 689
Meurs, E.J.A. & van den Heuvel, E.P.J. 1989, A&A, 226, 88
Miroshnichenko, A.S., Bjorkman, K.S., Chentsov, E.L. et al. 2002, A&A, 388, 563
Morgan, E., Remillard, R. & Swank, J. 2003, ATel 199
Motch, C., Haberl, F., Dennerl, K. et al. 1997, A&A, 323, 853
Negueruela, I. 2004, astro-ph/0411335
Negueruela, I., Smith, D.M., Reig, P. et al. 2005, astro-ph/0511088, in Proceedings of "The X-ray Universe 2005", held in San Lorenzo de El Escorial (Madrid), 26-30 September 2005, ESA-SP 604
Negueruela, I., Smith, D.M., Reig, P. et al. 2006, ApJ, 638, 982
Orosz, J.A., Kuulkers, E., van der Klis, M. et al. 2001, ApJ, 555, 489
Pakull, M.W., Motch, C. & Negueruela, I. 2003, ATel 202
Pellizza, L.J., Chaty, S., Negueruela, I. 2006, A&A, submitted
Pietsch, W., Mochejska, B.J., Misanovic, Z. et al. 2004, A&A, 413, 879
Reig, P. 1999, A&A, 345, 576
Remillard, R.A. & Smith, D.A. 2002, ATel 88
Ritter, H. & Kolb, U. 2003, A&A, 404, 301
Sakano, M., Koyama, K., Murakami, H. et al. 2002, ApJS, 138, 19
Skrutskie, M.F., Cutri, R.M.,Stiening, R. et al. 2006, AJ, 131, 1163.
Smith, D.M. et al. 2005, ApJ, 638, 974
Swank, J. & Morgan, E. 2000, IAUC 7531
Swartz, D.A., Ghosh, K.K., McCollough, M.L. et al. 2003, ApJS, 144, 213
Torii, K., Kinugasa, K., Katayama, K. et al. 1998, ApJ, 503, 843
Torrejon, J.M. & Orr, A. 2001, A&A, 377, 148
Walter, R. et al. 2004, HEAD 8, 3301
Winkler, C., Courvoisier, T.J.-L., Di Cocco, G. et al. 2003, A&A, 411, L1
Zezas, A., Fabbiano, G., Rots, A. H., Murray, S. S. 2002, ApJS, 142, 239
Zhang, F., Li, X.-D., Wang, Z.-R. 2004, ApJ, 603, 663

[1] Aab, O.E. et al. 1983, SvA 27, 603
[2] Aab, O.E. et al. 1984, SvAL, 9, 285
[3] Aab, O.E. et al. 1984, SvAL, 10, 386
[4] Abell, G.O. & Margon, B. 1979, Nat, 279, 701
[5] Aharonian, F. et al. 2005, Sci, 309, 746
[6] Anderson, S.F. et al. 1983, ApJ, 269, 605
[7] Anderson, S.F. et al. 1983, ApJ, 273, 697
[8] Angelini, L. et al. 1989, ApJ, 346, 906
[9] Angelini, L. et al. 1998, A&A, 339, L41
[10] Ankay, A. et al. 2001, A&A, 370, 170
[11] Aoki, T. et al. 1992, PASJ, 44, 461
[12] Apparao, K.M.V. et al. 1978, Nat, 271, 225
[13] Apparao, K.M.V. et al. 1980, A&A, 89, 249
[14] Armstrong, J.T. et al. 1980, ApJ, 236, L131
[15] Ash, T.D. et al. 1999, MNRAS, 307, 357
[16] Aslanov, A.A. et al. 1987, SvAL, 13, 369
[17] Audley, M.D. et al. 1996, ApJ, 457, 397
[18] Augello, G. et al. 2003, ApJ, 596, L63
[19] Balog, N.I. et al. 1983, SvA 27, 310
[20] Balona, L.A. et al. 1992, A&AS, 92, 533
[21] Balucinska-church, M. et al. 2000, MNRAS, 311, 861
[22] Bamba, A. et al. 2001, PASJ, 53, 1179
[23] Band, D.L. 1989, ApJ, 336, 937
[24] Band, D.L. & Gordon, M.A. 1989, ApJ, 338, 945
[25] Barnes, A.D. et al. 2005, MNRAS, astro-ph/0510448
[26] Barr, P. & Van der Woerd, H. 1990, ApJ, 352, L41
[27] Barr, P. et al. 1985, MNRAS, 216, 65P
[28] Barsukova, E.A. et al. 2005, ATel 416
[29] Barziv, O. et al. 2001, A&A, 377, 925
[30] Baykal et al. 2000, ApJ, 544, L129
[31] Becker, R.H. et al. 1977, ApJ, 216, L11
[32] Becklin, E.E. et al. 1972, Nat, 245, 302
[33] Beckmann, V. et al. 2005, ApJ, 631, 506
[34] Belloni, T. & Hasinger, G. 1990, A&A, 227, L33
[35] Belloni, T. & Hasinger, G. 1990, A&A, 230, 103
[36] Belloni, T. et al., 1993, A&A, 271, 487
[37] Belloni, T. et al. 1999, ApJ, 527, 345
[38] Berdnik, E.V. et al. 1990, SvAL, 16, 472
[39] Bergner, Yu.K. et al. 1995, A&AS, 112, 221
[40] Bernacca P.L. & Bianchi, L. 1981, A&A, 94, 345
[41] Beskrovnaya, N.G. 1989, SvAL, 14, 314
[42] Bianchi, L. & Bernacca P.L. 1980, A&A, 89, 214
[43] Bignami, G.F. et al. 1981, ApJ, 247, L85
[44] Bird, A. J. et al. 2004, ApJ, 607, L33
[45] Bisscheroux, B.C. et al. 1997, A&A, 317, 815
[46] Blay, P. et al. 2005, A&A, 438, 963
[47] Blay, P. et al. 2006, A&A, 446, 1095
[48] Blundell, K.M. et al. 2001, ApJ, 562, L79
[49] Blundell, K.M. & Bowler, M.G. 2004, ApJ, 616, L159
[50] Bodaghee, A. et al. 2004, ATel 290
[51] Bodaghee, A. et al. 2005, A&A, astro-ph/0510112
[52] Bolton, C.T. 1972, Nat, 235, 271
[53] Bonnet-Bidaud, J.M. & Mouchet, M. 1998, A&A, 332, L9
[54] Bonnet-Bidaud, J.M. & Van der Klis, M. 1981, A&A, 101, 299
[55] Bonning, E.W. & Falanga, M. 2005, A&A, 436, L31
[56] Bonsignori-Facondi, S.R. et al. 1986, A&A, 166, 157
[57] Bord, D.J. et al. 1976, ApJ, 203, 689
[58] Boroson, B. et al. 2003, ApJ, 592, 516
[59] Borozdin, K. et al. 1990, SvAL, 16, 345
[60] Bosch-Ramon, V. et al. 2005, ApJ, 628, 388
[61] Boyd, P.T. & Smale, A.P. 2004, ApJ, 612, 1006
[62] Boynton, P.E. et al. 1984, ApJ, 283, L53
[63] Boynton, P.E. et al. 1986, ApJ, 307, 545
[64] Bradt, H.V. & McClintock, J.E. 1983, ARA&A, 21, 63
[65] Bradt, H.V. et al. 1977, Nat, 269, 21
[66] Braes, L.L.E. & Miley, G.K. 1971, Nat, 232, 246
[67] Braes, L.L.E. & Miley, G.K. 1976, Nat, 264, 731
[68] Brinkmann, W. et al. 1989, A&A, 218, L13
[69] Brinkmann, W. et al. 1991, A&A, 241, 112
[70] Brucato, R.J. et al. 1972, ApJ, 175, L137
[71] Bruhweiler, F.C. et al. 1981, ApJS, 46, 255
[72] Burderi, L. et al. 2000, ApJ, 530, 429
[73] Buscombe, W. 1962, MNRAS, 124, 189
[74] Buscombe, W. 1980, MK Classification, 4th General Catalogue (Evanston)
[75] Cameron, P.B. et al. 2004, ATel 312
[76] Campana, S. et al. 1999, A&A, 352, L91
[77] Campana, S. et al. 2001, ApJ, 561, 924
[78] Carpenter, G.F. et al. 1977, MNRAS, 179, 27P

[79] Casares, J. et al. 2005, MNRAS, 360, 1105
[80] Casares, J. et al. 2005, MNRAS, 364, 899
[81] Chakrabarty, D. et al. 1993, ApJ, 403, L33
[82] Chakrabarty, D. et al. 1995, ApJ, 446, 826
[83] Chakrabarty, D. et al. 2002, ApJ, 573, 789
[84] Chakrabarti, S.K. et al. 2003, ApJ, 595, L45
[85] Chakrabarti, S.K. et al. 2005, MNRAS, 362, 957
[86] Charles, P.A. et al. 1975, Ap. Lett. 16, 145
[87] Chaty, S. et al. 2003, MNRAS, 343, 169
[88] Chenevez, J. et al. 2004, ATel 223
[89] Cherepashchuk, A.M. et al. 2003, ATel 159
[90] Cherepashchuk, A.M. et al. 2005, A&A, 437, 561
[91] Chernyakova, M. et al. 2003, ATel 157
[92] Chernyakova, M. et al. 2005, ATel 519
[93] Chernyakova, M. et al. 2005, MNRAS, astro-ph/0508515
[94] Chevalier, C. & Ilovaisky, S.A. 1998, A&A, 330, 201
[95] Choudhury, M. & Rao, A.R. 2004, ApJ, 616, L143
[96] Cieslinski, D. et al. 1994, A&AS, 106, 243
[97] Clark, G.W. 1975, IAU Circ 2843
[98] Clark, G.W. 1990, ApJ, 353, 274
[99] Clark, G.W. 2000, ApJ, 542, L131
[100] Clark, G.W. 2004, ApJ, 610, 956
[101] Clark, G.W. et al. 1988, ApJ, 324, 974
[102] Clark, J.S. et al. 1998, MNRAS, 294, 165
[103] Clark, J.S. et al. 1998, MNRAS, 297, 657
[104] Clark, J.S. et al. 1999, A&A, 348, 888
[105] Clark, J.S. et al. 2000, A&A, 356, 50
[106] Clark, J.S. et al. 2002, A&A, 392, 909
[107] Clarke, D. 1990, A&A, 227, 151
[108] Coburn, W. 2005, ATel 381
[109] Coburn, W. et al. 2001, ApJ, 552, 738
[110] Coburn, W. et al. 2004, ATel 337
[111] Codina, S.J. et al. 1984, A&AS, 57, 239
[112] Coe, M.J. & Payne, B.J. 1985, Ap&SS 109, 175
[113] Coe, M.J. et al. 1987, MNRAS, 226, 455
[114] Coe, M.J. et al. 1988, MNRAS, 232, 865
[115] Coe, M.J. et al. 1990, MNRAS, 243, 475
[116] Coe, M.J. et al. 1993, MNRAS, 261, 599
[117] Coe, M.J. et al. 1994, MNRAS, 270, L57
[118] Coe, M.J. et al. 1994, A&A, 289, 784
[119] Coe, M.J. et al. 1996, MNRAS, 281, 333
[120] Combi, J.A. et al. 2004, A&A, 422, 1031
[121] Combi, J.A. et al. 2004, A&A, 427, 959
[122] Cominsky, L. & Moraes, F. 1991, ApJ, 370, 670
[123] Cominsky, L. et al. 1978, ApJ, 224, 46
[124] Conti, P.S. 1978, A&A, 63, 225
[125] Conti, P.S. & Cowley, A.P. 1975, ApJ, 200, 133
[126] Cook, M.C. et al. 1984, in "X-ray Astronomy '84", eds. M. Oda & R. Giacconi, p. 225
[127] Cook, M.C. & Page, C.G. 1987, MNRAS, 225, 381
[128] Cook, M.C. & Warwick, R.S. 1987, MNRAS, 225, 369; 227, 661
[129] Cooke, B.A. et al. 1984, ApJ, 285, 258
[130] Corbet, R.H.D. & Day, C.S.R. 1990, MNRAS, 243, 553
[131] Corbet, R.H.D. & Mason, K.O. 1984, A&A, 131, 355
[132] Corbet, R.H.D. & Mukai, K. 2002, ApJ, 577, 923
[133] Corbet, R.H.D. & Peele, A.G. 2000, ApJ, 530, L33
[134] Corbet, R.H.D. & Peele, A.G. 2001, ApJ, 562, 936
[135] Corbet, R.H.D. & Remillard, R. 2005, ATel 377
[136] Corbet, R.H.D. et al. 1986, MNRAS, 221, 961
[137] Corbet, R.H.D. et al. 1986, A&A, 162, 117
[138] Corbet, R.H.D. et al. 1997, ApJ, 489, L83
[139] Corbet, R.H.D. et al. 1999, ApJ, 517, 956
[140] Corbet, R.H.D. et al. 2004, ATel 269
[141] Corbet, R.H.D. et al. 2005, ApJ, 633, 377
[142] Corbet, R.H.D. et al. 2005, ATel 649
[143] Courvoisier, T. J.-L. et al. 2003, IAUC, 8063
[144] Cox, N.L.J. et al. 2005, A&A, 436, 661
[145] Coyne, G.V. & McConnell, D.J. 1983, Vatican Obs. Publ. 1, part no 6, 73
[146] Crampton, D. & Hutchings, J.B. 1981, ApJ, 251, 604
[147] Crampton, D. et al. 1978, ApJ, 225, L63
[148] Crampton, D. et al. 1985, ApJ, 299, 839
[149] Crawford, I.A. 1991, A&A, 246, 210
[150] Cusumano, G. et al. 1998, A&A, 338, L79
[151] Cusumano, G. et al. 2000, ApJ, 528, L25
[152] Dachs, J. 1970, A&A, 5, 312
[153] Dachs, J. 1972, A&A, 21, 373
[154] Dachs, J. 1976, A&A, 47, 19
[155] Dachs, J. et al. 1989, A&AS, 78, 487
[156] D'Amico, N. et al. 1987, A&A, 180, 114
[157] Davis, R. & Hartmann, L. 1983, ApJ, 270, 671
[158] Davison, P.J.N. et al. 1977, MNRAS, 181, 73P
[159] Deeter, J.E. et al. 1987, ApJ, 314, 634
[160] Deeter, J.E. et al. 1987, AJ, 93, 877
[161] Deeter, J.E. et al. 1989, ApJ, 336, 376
[162] Delgado-Marti, H. et al. 2001, ApJ, 546, 455
[163] De Loore, C. et al. 1981, A&A, 104, 150
[164] De Loore, C. et al. 1984, A&A, 141, 279
[165] den Hartog, P.R. et al. 2004, ATel 281
[166] Densham, R.H. & Charles, P.A. 1982, MNRAS, 201, 171
[167] Dickey, J.M. 1983, ApJ, 273, L71
[168] Di Salvo, T. et al. 2001, ApJ, 547, 1024
[169] Doazan, V. et al. 1987, A&A, 182, L25
[170] D'Odorico, S. et al. 1991, Nat, 353, 329
[171] Dolan, J.F. & Tapia, S. 1984, A&A, 139, 249
[172] Dolan, J.F. & Tapia, S. 1988, A&A, 202, 124
[173] Dolan, J.F. & Tapia, S. 1989, ApJ, 344, 830
[174] Doll, H. & Brinkmann, W. 1987, A&A, 173, 86
[175] Dower, R.G. et al. 1978, Nat, 273, 364
[176] Downes, R.A. 1984, PASP, 96, 807
[177] Doxsey, R.E. et al. 1977, Nat, 269, 112
[178] Drake, S.A. 1990, AJ, 100, 572
[179] Drilling, J.S. 1973, AJ, 80, 128
[180] Drilling, J. S. 1991, ApJS, 76, 1033
[181] Dupree, A. et al. 1978, Nat, 275, 400
[182] Dupree, A. et al. 1980, ApJ, 238, 969
[183] Eikenberry, S.S. et al. 2001, ApJ, 561, 1027
[184] Estalella, R.A. et al. 1993, A&A, 268, 178
[185] Exosat & Einstein Data Bases
[186] Eyles, C.J. et al. 1975, Nat, 254, 577
[187] Fabbiano, G. & Schreier, E.J. 1977, ApJ, 214, 235
[188] Fabian, A.C. et al. 1989, MNRAS, 238, 729
[189] Fabregat, J. et al. 1992, A&A, 259, 522
[190] Fabrika, S.N. & Bychkova, L.V. 1990, A&A, 240, L5
[191] Fahlman, G.G. & Walker, G.A.H. 1980, ApJ, 240, 169
[192] Fahlman, G.G. et al. 1985, ApJS, 58, 225
[193] Falomo, R. et al. 1987, MNRAS, 224, 323
[194] Farrell, S.A. et al. 2005, MNRAS, astro-ph/0502008
[195] Fejes, L. 1986, A&A, 168, 69
[196] Fejes, L. et al. 1988, A&A, 189, 124
[197] Fender, R.P. & Hendry, M.A. 2000, MNRAS, 317, 1
[198] Fender, R. et al. 1999, MNRAS, 308, 473
[199] Fender, R. et al. 2000, ApJ, 530, L29
[200] Fender, R.P. et al. 2000, MNRAS, 312, 853
[201] Feng, H. et al. 2004, ApJ, 612, L45
[202] Ferrer, O.E. 1980, A&A, 84, 108
[203] Fiedler, R.L. et al. 1987, AJ, 94, 1244
[204] Filippenko, A.V. et al. 1988, AJ, 96, 242
[205] Filliatre, P. & Chaty, S. 2004, ApJ, 616, 469
[206] Finger, M.H. et al. 1994, IAU Circ 5959
[207] Finger, M.H. et al. 1996, A&AS, 120, 209
[208] Finger, M.H. et al. 1996, ApJ, 459, 288
[209] Finger, M.H. et al. 1999, ApJ, 517, 449
[210] Finley, J.P. et al. 1992, A&A, 262, L25
[211] Finley, J.P. et al. 1994, ApJ, 429, 356
[212] Fitzgerald, M.P. 1973, A&AS, 9, 297
[213] Forman, W. et al. 1973, ApJ, 182, L103
[214] Forman, W. et al. 1976, ApJ, 206, L29
[215] Forman, W. et al. 1978, ApJS, 38, 357
[216] Foschini, L. et al. 2003, IAUC, 8076
[217] Frail, D.A. & Hjellming, R.M. 1991, AJ, 101, 2126
[218] Friend, D.B. & Cassinelli, J.P. 1986, ApJ, 303, 292
[219] Frontera, F. & Fuligni, F. 1975, ApJ, 198, L105
[220] Frontera, F. et al. 1987, ApJ, 320, L127
[221] Frontera, F. et al. 1998, A&A, 339, L69
[222] Gallo, E. et al. 2005, Nat, 436, 819
[223] Galloway, D.K. et al. 2004, ApJ, 613, 1164
[224] Galloway, D.K. et al. 2005, ApJ, astro-ph/0506247
[225] Geldzahler, B.J. et al. 1983, ApJ, 273, L65
[226] Geldzahler, B.J. et al. 1989, ApJ, 342, 1123
[227] Giacconi, R. et al. 1971, ApJ, 167, L67
[228] Giangrande, A. et al. 1980, A&AS, 40, 289
[229] Gierlinski, M. & Zdziarski, A.A. 2003, MNRAS, 343, L84
[230] Gierlinski, M. et al. 1997, MNRAS, 288, 958
[231] Gies, D.R. & Bolton, C.T. 1982, ApJ, 260, 240
[232] Gies, D.R. & Bolton, C.T. 1984, ApJ, 276, L17
[233] Gies, D.R. & Bolton, C.T. 1986, ApJ, 304, 371

[234] Gies, D.R. & Bolton, C.T. 1986, ApJ, 304, 389
[235] Gies, D. R. et al. 2002, ApJ, 566, 1069
[236] Gies, D. R. et al. 2003, ApJ, 583, 424
[237] Gilfanov, M. et al. 1999, A&A, 352, 182
[238] Glass, I.S. 1979, MNRAS, 187, 807
[239] Gleissner, T. et al. 2004, A&A, 425, 106
[240] Gogus, E. et al. 2005, ApJ, 632, 1069
[241] Goldstein, G. et al. 2004, ApJ, 127, 2310
[242] Golenetskii, S. et al. 2003, ApJ, 596, 1113
[243] Gonzalez-Riestra, R. et al. 2004, A&A, 420, 589
[244] Goranskij, V.P. 1990, IBVS 3464, 1
[245] Goranskij, V.P. et al. 1991, SvAL, 17, 399
[246] Gottwald, M. et al. 1986, MNRAS, 222, 21P
[247] Gottwald, M. et al. 1991, A&AS, 89, 367
[248] Grankin, K.N. et al. 1991, SvAL, 17, 415
[249] Grebenev, S. & Sunyaev, R. 1991, IAU Circ 5354
[250] Grebenev, S.A. et al. 2003, ATel 252
[251] Grebenev, S.A. et al. 2004, ATel 275
[252] Grebenev, S. A. et al. 2004, ATel 350
[253] Grebenev, S.A. et al. 2005, ATel 457
[254] Gregory, P.C. 2002, ApJ, 575, 427
[255] Gregory, P.C. et al. 1972, Nat, 239, 440
[256] Gregory, P.C. et al. 1979, AJ, 84, 1030
[257] Grillo, E. et al. 1992, ApJS, 81, 795
[258] Grindlay, J.E. et al. 1984, ApJ, 276, 621
[259] Grindlay, J.E. et al. 1984, ApJ, 277, 286
[260] Grove, J.E. et al. 1995, ApJ, 438, L25
[261] Gruber, D.E. 1988, ApJ, 328, 265
[262] Haberl, F. 1991, ApJ, 376, 245
[263] Haberl, F. 1995, A&A, 296, 685
[264] Haberl, F. & White, N.E. 1990, ApJ, 361, 225
[265] Haberl, F. et al. 1989, ApJ, 343, 409
[266] Haberl, F. et al. 1998, A&A, 335, 587
[267] Haigh, N.J. et al. 1999, MNRAS, 310, L21
[268] Halpern, J.P., Gotthelf, E.V. 2004, ATel 319
[269] Halpern, J.P., Gotthelf, E.V. 2004, ATel 344
[270] Halpern, J.P. & Tyagi, S. 2005, ATel, 681
[271] Halpern, J.P. et al. 2004, ATel 289
[272] Halpern, J.P. & Tyagi, S. 2005, ATel 682
[273] Hammerschlag-Hensberge, G. 1978, A&A, 64, 399
[274] Hammerschlag-Hensberge, G. et al. 1979, A&A, 76, 245
[275] Hammerschlag-Hensberge, G. et al. 1980, A&A, 85, 119
[276] Hammerschlag-Hensberge, G. et al. 1990, ApJ, 352, 698
[277] Hammerschlag-Hensberge, G. et al. 2003, A&A, 407, 685
[278] Hardorp, J. et al. 1959, Luminous Stars in the Northern Milky Way I (Hamberger Sternwarte- Warner and Swasey Observatory)
[279] Harmanec, P. et al. 2000, A&A, 364, L85
[280] Harrison, T. et al. 2000, ApJ, 528, 454
[281] Heap, S. & Corcoran, M.F. 1992, ApJ, 387, 340
[282] Heindl, W.A. et al. 1999, ApJ, 521, L48
[283] Heindl, W.A. et al. 2001, ApJ, 563, L35
[284] Heindl, W.A. et al. 2003, ApJ, 588, L97
[285] Heindl, W.A. et al. 2003, ATel 200
[286] Henrichs, H. et al. 1982, ApJ, 268, 807
[287] Hensberge, G. et al. 1973, A&A, 29, 69
[288] Henson, G.D. et al. 1983, ApJ, 275, 247
[289] Hermsen, W. et al. 1977, Nat, 269, 494
[290] Herrero, A. et al. 1995, A&A, 297, 556
[291] Hill, A.B. et al. 2005, A&A, 439, 255
[292] Hill, R. et al. 1974, ApJ, 189, L69
[293] Hillwig, T.C. et al. 2004, ApJ, 615, 422
[294] Hjellming, R.M. 1973, ApJ, 182, L29
[295] Hjellming, R.M. 1997, IAU Circ. 6750
[296] Hjellming, R.M. & Johnston, K.J. 1981, ApJ, 246, L141
[297] Hjellming, R.M. & Wade, C.M. 1971, ApJ, 168, L21
[298] Hjellming, R.M. et al. 2000, ApJ, 544, 977
[299] Hoffleit, D. & Jaschek, C. 1982, Bright Star Catalogue (Yale)
[300] Hog, E. et al. 1998, A&A, 335, L65
[301] Holt, S.S. et al. 1979, ApJ, 233, 344
[302] Honeycutt, R.K. & Schlegel, E.M. 1985, PASP, 97, 300
[303] Hong, J. & Hailey, C.J. 2004, ApJ, 600, 743
[304] Hony, S. et al. 2000, A&A, 355, 187
[305] Horaguchi, T. et al. 1994, PASJ, 46, 9
[306] Howarth, I.D. 1983, MNRAS, 203, 801
[307] Howe, S.K. et al. 1983, ApJ, 272, 678
[308] Howells, L. et al. 2001, A&A, 369, 99
[309] Huckle, H.E. et al. 1977, MNRAS, 180, 21P
[310] Hulleman, F. et al. 1998, A&A, 337, L25
[311] Hunt, L.K. et al. 1994, A&A, 290, 428
[312] Hutchings, J.B. 1970, MNRAS, 150, 55
[313] Hutchings, J.B. 1974, ApJ, 192, 677
[314] Hutchings, J.B. 1977, MNRAS, 181, 619
[315] Hutchings, J.B. 1979, PASP, 91, 657
[316] Hutchings, J.B. & Crampton, D. 1981, PASP, 93, 486
[317] Hutchings, J.B. & Crampton, D. 1981, ApJ, 247, 222
[318] Hutchings, J.B. & Stoeckley, T.R. 1977, PASP, 89, 19
[319] Hutchings, J.B. et al. 1975, MNRAS, 170, 313
[320] Hutchings, J.B. et al. 1978, ApJ, 223, 530
[321] Hutchings, J.B. et al. 1979, ApJ, 229, 1079
[322] Hutchings, J.B. et al. 1981, AJ, 86, 871
[323] Hutchings, J.B. et al. 1982, PASP, 94, 541
[324] Hutchings, J.B. et al. 1984, PASP, 96, 312
[325] Hutchings, J.B. et al. 1987, PASP, 99, 420
[326] Huthoff, F. & Kaper, L. 2002, A&A, 383, 999
[327] Hynes, R. I. et al. 2002, A&A, 392, 991
[328] Iaria, R. et al. 2005, ApJ, 634, L161
[329] Ibragimov, A. et al. 2005, MNRAS, 362, 1435
[330] Ilovaisky, S.A. et al. 1979, A&A, 71, L17
[331] Ilovaisky, S.A. et al. 1982, A&A, 114, L7
[332] Inam, S.C. et al. 2004, ApJ, 616, 463
[333] Inam, S. C. et al. 2004, MNRAS, 349, 173
[334] in't Zand, J.J.M. 2005, A&A, 441, L1
[335] in't Zand, J.J.M. & Heise, J. 2004, ATel 362
[336] in't Zand, J.J.M. et al. 1997, ApJ, 479, L47
[337] in't Zand, J.J.M. et al. 1998, ApJ, 496, 386
[338] in't Zand, J.J.M. et al. 1998, IAUC 6840
[339] in't Zand, J.J.M. et al. 2000, A&A, 357, 520
[340] in't Zand, J.J.M. et al. 2000, A&A, 361, 85
[341] in't Zand, J.J.M. et al. 2001, A&A, 380, L26
[342] in't Zand, J.J.M. et al. 2001, ApJ, 553, L165
[343] in't Zand, J.J.M. et al. 2004, Proceedings of the 5th INTEGRAL Workshop on the INTEGRAL Universe (ESA SP-552), eds.: V. Schönfelder, G. Lichti & C. Winkler, 16-20 February 2004, Munich, Germany, p.427. astro-ph/0407085
[344] in't Zand, J.J.M. et al. 2005, astro-ph/0505256
[345] Iping, R.C. & Petterson, J.A. 1990, A&A, 239, 221
[346] Ishida, M. et al. 2004, ApJ, 601, 1088
[347] Israel, G.L. et al. 1997, ApJ, 474, L53
[348] Israel, G.L. et al. 1998, MNRAS, 298, 502
[349] Israel, G.L. et al. 2000, MNRAS, 314, 87
[350] Israel, G.L. et al. 2000, Proceedings of the 175th IAU Colloq., eds. M.A. Smith, H.F. Henrichs, & J. Fabregat, ASP Conf. Ser. 214, 739
[351] Israel, G.L. et al. 2001, A&A, 371, 1018
[352] Ives, J.C. et al. 1975, Nat, 254, 578
[353] Iye, M. 1986, PASP, 38, 463
[354] Iye, M. & Kodaiza, K. 1986, PASP, 97, 1186
[355] Janot-Pacheco, E. et al. 1981, A&A, 99, 274
[356] Janot-Pacheco, E. et al. 1987, A&A, 177, 91
[357] Janot-Pacheco, E. et al. 1988, A&A, 202, 81
[358] Jarad, M.M. et al. 1989, MNRAS, 230, 1085
[359] Jaschek, M. & Jaschek, C. 1963, PASP, 73, 365
[360] Johnston, K.J. et al. 1986, ApJ, 309, 707
[361] Johnston, M. et al. 1978, ApJ, 223, L71
[362] Jones, C. et al. 1973, ApJ, 181, L43
[363] Kaaret, P. et al. 1999, ApJ, 523, 197
[364] Kaaret, P. et al. 2000, ApJ, 542, L41
[365] Kallman, T.R. et al. 1987, ApJ, 317, 746
[366] Kamata, Y. et al. 1990, PASJ, 42, 785
[367] Kaper, L. et al. 1990, Nat, 347, 652
[368] Kaper, L. et al. 1995, A&A, 300, 446
[369] Kaper, L. et al. 1997, ApJ, 475, L37
[370] Kato, T. et al. 1999, IBVS 4777
[371] Katz, J.L. et al. 1984, AJ, 89, 1604
[372] Kawai, N. et al. 1989, PASJ, 41, 491
[373] Kelley, R.L. et al. 1981, ApJ, 243, 251
[374] Kelley, R.L. et al. 1983, ApJ, 268, 790
[375] Kelley, R.L. et al. 1983, ApJ, 274, 765
[376] Kemp, J.C. et al. 1983, ApJ, 271, L65
[377] Kemp, J.C. et al. 1986, ApJ, 305, 805
[378] Kemp, J.C. et al. 1987, SvA 31, 170
[379] Kennea, J. A. et al. 2005, ATel 599
[380] Kennea, J. A. et al. 2005, ATel 673
[381] King, A.R. et al. 2000, ApJ, 530, L25
[382] Kinugaso, K. et al. 1998, ApJ, 495, 435
[383] Kitamoto, S. et al. 1987, PASJ, 39, 259
[384] Kitamoto, S. et al. 1989, PASJ, 41, 81

[385] Kitamoto, S. et al. 1990, PASJ, 42, 85
[386] Kitamoto, S. et al. 1992, ApJ, 384, 263
[387] Klare, G. & Neckel, T. 1977, A&AS, 27, 215
[388] Kniffen, D.A. et al. 1997, ApJ, 486, 126
[389] Kodaira, K. et al. 1985, PASJ, 37, 97
[390] Kodaira, K. et al. 1985, ApJ, 296, 232
[391] Koenigsberger, G. et al. 1983, ApJ, 273, 716
[392] Koenigsberger, G. et al. 2003, RMxAA, 39, 17
[393] Koh, D.T. et al. 1997, ApJ, 479, 933
[394] Kohoutek, L., Wehmeyer, R. 1999, A&AS, 134, 255
[395] Konig, M. & Maisack, M. 1997, ApJ, 327, L33
[396] Kopylov, I.M. et al. 1986, SvA 30, 408
[397] Kopylov, I.M. et al. 1987, SvA 31, 410
[398] Kotani, T. et al. 2005, astro-ph/0509411
[399] Koyama, K. et al. 1990, Nat, 343, 148
[400] Koyama, K. et al. 1990, ApJ, 356, L47
[401] Koyama, K. et al. 1990, PASJ, 42, L59
[402] Koyama, K. et al. 1991, ApJ, 366, L19
[403] Koyama, K. et al. 1991, ApJ, 370, L77
[404] Kretschmar, P. et al. 2005, ATel 601
[405] Kreykenbohm, I. et al. 2002, A&A, 395, 129
[406] Kreykenbohm, I. et al. 2004, A&A, 427, 975
[407] Kriss, G.A. et al. 1983, ApJ, 266, 806
[408] Krzeminsky, W. et al. 1974, ApJ, 192, L135
[409] Kunjaya, C. & Hirata, R. 1995, PASJ, 47, 589
[410] Kuulkers, E. 2005, astro-ph/0504625
[411] Kuulkers, E. et al. 2003, ATel 149
[412] La Barbera, A. et al. 2003, A&A, 400, 993
[413] La Barbera, A. et al. 2005, A&A, 438, 617
[414] Lamb, R.C. et al. 1980, ApJ, 239, 651
[415] Lamb, R.C. et al. 1983, Nat, 305, 37
[416] Lapshov, I. et al. 1992, SvAL, 18, 12
[417] Laycock, S. et al. 2003, MNRAS, 338, 211
[418] Leahy, D.A. 2001, A&A, 380, 516
[419] Leahy, D.A. 2002, A&A, 391, 219
[420] Leahy, D.A. & Ananth, A.G. 1992, MNRAS, 256, 39P
[421] Leahy, D.A. & Matsoka, M. 1990, ApJ, 355, 627
[422] Leahy, D.A. et al. 1988, PASJ, 40, 197
[423] Leahy, D.A. et al. 1989, MNRAS, 237, 269
[424] Leibowitz, E.M. 1984, MNRAS, 210, 279
[425] Leibowitz, E.M. & Mendelson, H. 1992, PASP, 94, 977
[426] Leibowitz, E.M. et al. 1984, MNRAS, 206, 751
[427] Leibowitz, E.M. et al. 1984, Nat, 307, 341
[428] Levine, A.M. et al. 2004, ApJ, 617, 1284
[429] Li, F. et al. 1979, ApJ, 228, 893
[430] Li, X.D. & van den Heuvel, E.P.J. 1999, ApJ, 513, L45
[431] Ling, J.C. et al. 1983, ApJ, 275, 307
[432] Ling, J.C. et al. 1987, ApJ, 321, L117
[433] Ling, J.C. et al. 1997, ApJ, 484, 375
[434] Liu, Q.Z. & Hang, H.R. 1999, A&A, 350, 855
[435] Liu, Q.Z. & Hang, H.R. 2001, Ap&SS, 275, 401
[436] Liu, Q.Z. & Mirabel, I.F. 2005, A&A, 429, 1125
[437] Liu, Q.Z. & Yan, J.Z. 2005, NewA, 11, 130
[438] Liu, Q.Z. et al. 2000, A&A, 359, 646
[439] Liu, Q.Z. et al. 2000, A&AS, 147, 25 (3rd edition)
[440] Liu, Q.Z. et al. 2005, A&A, 442, 1135
[441] Lochner, J.C. et al. 1989, ApJ, 337, 823
[442] Lochner, J.C. et al. 1991, ApJ, 376, 295
[443] Lu, F.J. et al. 1996, A&AS, 115, L395
[444] Lubinski, P. et al. 2005, ATel 469
[445] Lum, K.S.K. et al. 1992, ApJS, 78, 423
[446] Lutovinov, A. 2004, ATel 329
[447] Lutovinov, A. et al. 2005, A&A, 444, 821
[448] Lutovinov, A. & Revnivtsev, M. 2003, AstL, 29, 719
[449] Lutovinov, A. et al. 2003, ATel 155
[450] Lutovinov, A. et al. 2005, A&A, 433, L41
[451] Lyubimkov, L.S. et al. 1997, MNRAS, 286, 549
[452] Lyuty, V.M. & Zaitseva, G.V. 2000, AstL, 26, 9
[453] Lyutyi, V.M. 1985, SvA 29, 429
[454] Lyutyi, V.M. et al. 1989, SvAL, 15, 182
[455] Maccarone, T.J. & Coppi, P.S. 2002, MNRAS, 335, 465
[456] MacConnell, D.J. 1982, A&AS, 48, 355
[457] Maitra, D. & Bailyn, C. D. 2006, ApJ, 637 in press
[458] Makishima, K. et al. 1984, PASJ, 36, 679
[459] Makishima, K. et al. 1987, ApJ, 314, 619
[460] Makishima, K. et al. 1990, ApJ, 365, L59
[461] Makishima, K. et al. 1990, PASP, 42, 295
[462] Malizia, A. et al. 2005, ApJ, 630, L157
[463] Maraschi, L. et al. 1981, ApJ, 248, 1010
[464] Margon, B. & Anderson, S.F. 1989, ApJ, 347, 448
[465] Margon, B. et al. 1977, ApJ, 216, 811
[466] Margon, B. et al. 1979, ApJ, 230, L41
[467] Margon, B. et al. 1979, ApJ, 233, L63
[468] Margon, B. et al. 1984, ApJ, 281, 313
[469] Markert, T.H. et al. 1978, ApJ, 224, 157
[470] Markert, T.H. et al. 1979, ApJS, 39, 573
[471] Markwardt, C.B. et al. 2004, ATel 317
[472] Markwardt, C.B. et al. 2005, ATel 465
[473] Marsden, D. et al. 1998, ApJ, 502, L129
[474] Marshall, F. E. et al. 1999, IAUC, 7240
[475] Marshall, F.E. et al. 2000, IAUC, 7363
[476] Marshall, H.L. et al. 2002, ApJ, 564, 941
[477] Marshall, N. & Ricketts, M.J. 1980, MNRAS, 193, 7P
[478] Marti, J. & Paredes, J. M. 1995, A&A, 298, 151
[479] Marti, J. et al. 1998, A&A, 338, L71
[480] Marti, J. et al. 2000, ApJ, 545, 939
[481] Marti, J. et al. 2001, A&A, 375, 476
[482] Marti, J. et al. 2004, A&A, 418, 271
[483] Marti, J. et al. 2005, A&A, 439, 279
[484] Martocchia, A. et al. 2005, A&A, 430, 245
[485] Masetti, N. et al. 2004, A&A, 423, 311
[486] Masetti, N. et al. 2005, astro-ph/0508451
[487] Masetti, N. et al. 2005, ATel 528
[488] Mason, K.O. et al. 1976, ApJ, 207, 78
[489] Mason, K.O. et al. 1978, MNRAS, 184, 45P
[490] Mason, K.O. et al. 1986, ApJ, 309, 700
[491] Massay, P. et al. 1995, ApJ, 454, 151
[492] Massi, M. 2004, A&A, 422, 267
[493] Massi, M. et al. 1993, A&A, 269, 249
[494] Massi, M. et al. 2001, A&A, 376, 217
[495] Massi, M. et al. 2004, A&A, 414, L1
[496] Massi, M. et al. 2005, AIPC, 745, 311
[497] Matsuoka, M. et al. 1986, MNRAS, 222, 605
[498] Matt, G. & Guainazzi, M. 2003, MNRAS, 341, L13
[499] Mazeh, T. et al. 1987, ApJ, 317, 824
[500] McCollough, M.L. et al. 1999, ApJ, 517, 951
[501] McConnell, M.L. et al. 2002, ApJ, 572, 984
[502] McCray, R. et al. 1984, ApJ, 282, 245
[503] McSwain, M.V. & Gies, D. R. 2002, ApJ, 568, L27
[504] McSwain, M.V. et al. 2004, ApJ, 600, 927
[505] Meekins, J.F. et al. 1984, ApJ, 278, 288
[506] Mendelson, H. & Mazeh, T. 1989, MNRAS, 239, 733
[507] Mendelson, H. & Mazeh, T. 1991, MNRAS, 250, 373
[508] Mereghetti, S. et al. 1987, ApJ, 312, 755
[509] Mereghetti, S. et al. 1991, ApJ, 366, L23
[510] Mereghetti, S. et al. 2000, A&A, 354, 567
[511] Migliari, S. et al. 2002, Sci, 297, 1673
[512] Mihara, T. 1995, Ph.D. thesis, Univ. of Tokyo
[513] Mihara, T. et al. 1991, ApJ, 379, L61
[514] Mikkelsen, D.R. & Wallerstein, G. 1974, ApJ, 194, 459
[515] Miller J.M. et al., 2002, ApJ, 577, L15
[516] Miller, J.M. et al. 2002, ApJ, 578, 348
[517] Miller, J.M. et al. 2005, ApJ, 620, 398
[518] Miller-Jones, J.C.A. et al. 2004, ApJ, 600, 368
[519] Mioduszewski, A.J. & Rupen, M.P. 2004, ApJ, 615, 432
[520] Mioduszewski, A.J. et al. 2001, ApJ, 553, 766
[521] Mirabel, I.F. & Rodrigues, I. 2003, Sci, 300, 1119
[522] Mirabel, I.F. et al. 2004, A&A, 422, L29
[523] Miroshnichenko, A.S. et al. 2002, A&A, 390, 627
[524] Mitani, K. et al. 1984, Ap&SS 103, 345
[525] Miyamoto, S. & Kitamoto, S. 1989, Nat, 342, 773
[526] Miyamoto, S. et al. 1989, Nat, 336, 450
[527] Miyamoto, S. et al. 1992, ApJ, 391, L21
[528] Moffat, A.F.J. et al. 1973, A&A, 23, 433
[529] Molkov, S.V. et al. 2004, ATel 274
[530] Molnar, L.A. & Mauche, C.W. 1986, ApJ, 310, 343
[531] Molnar, L.A. et al. 1984, Nat, 310, 662
[532] Mook, D.E. et al. 1974, PASP, 86, 894
[533] Morel, T. & Grosdidier, Y. 2005, MNRAS, 356, 665
[534] Morgan, E. et al. 2003, ATel 199
[535] Motch, C. & Janot-Pacheco, E. 1987, A&A, 182, L55
[536] Motch, C. et al. 1988, A&A, 201, 63
[537] Motch, C. et al. 1991, ApJ, 369, 490
[538] Motch, C. et al. 1991, A&A, 246, L24
[539] Motch, C. et al. 1997, A&A, 323, 853
[540] Motch, C. et al. 1998, A&AS, 132, 341

[541] Mouchet, M. et al. 1980, A&A, 90, 113
[542] Mourard, D. et al. 1989, Nat, 342, 520
[543] Murakami, H. et al. 2003, IAUC, 8070
[544] Murakami, T. et al. 1983, ApJ, 264, 563
[545] Murakami, T. et al. 1986, ApJ, 310, L31
[546] Murdin, P. et al. 1980, MNRAS, 193, 135
[547] Nagase, F. 1989, PASJ, 41, 1
[548] Nagase, F. et al. 1986, PASJ, 38, 547
[549] Nagase, F. et al. 1991, ApJ, 375, L49
[550] Nagase, F. et al. 1992, ApJ, 396, 147
[551] Namiki, M. et al. 2003, PASJ, 55, 281
[552] Negueruela, I. 1998, A&A, 338, 505
[553] Negueruela, I. et al. 1999, MNRAS, 307, 695
[554] Negueruela, I. & Okazaki, A.T. 2001, A&A, 369, 108
[555] Negueruela, I. & Reig, P. 2001, A&A, 371, 1056
[556] Negueruela, I. et al. 1996, A&A, 315, 160
[557] Negueruela, I. et al. 1997, MNRAS, 284, 859
[558] Negueruela, I. et al. 2000, A&A, 356, 1003
[559] Negueruela, I. et al. 2003, A&A, 397, 739
[560] Negueruela, I. et al. 2005, ATel 429 (see astro-ph/0511088)
[561] Negueruela, I. et al. 2005, ATel 470
[562] Negueruela, I. et al. 2006, ApJ, 638, 982
[563] Neshpor, Yu. I. et al. 2003, AstL, 29, 236
[564] Nicolet, B. 1978, A&AS, 34, 1
[565] Ninkov, Z. et al. 1987, ApJ, 321, 425 & 438
[566] Norton, A.J. et al. 1991, MNRAS, 253, 579
[567] Ogley, R.N. et al. 2001, MNRAS, 326, 349O
[568] Okazaki, A. T. & Negueruela, I. 2001, A&A, 377, 161
[569] Oosterbrook, T. et al. 1999, A&A, 351, L33
[570] Orlandini, M. et al. 2000, AdSpR, 25, 417
[571] Orlandini, M. et al. 2000, A&A, 356, 163
[572] Orosz J.A. et al. 2001, ApJ, 555, 489
[573] Osmer, P.S. et al. 1975, ApJ, 195, 705.
[574] Paerels, F. et al. 2000, ApJ, 533, L135
[575] Pakull, M. et al. 1983, A&A, 122, 79
[576] Pakull, M.W. et al. 2003, ATel 202
[577] Pandey, M. et al. 2005, A&A, astro-ph/0509645
[578] Paredes, J.M. & Figueras, F. 1986, A&A, 154, L30
[579] Paredes, J.M. et al. 1990, A&A, 232, 377
[580] Paredes, J.M. et al. 1994, A&A, 288, 519
[581] Paredes, J.M. et al. 1997, A&A, 320, L25
[582] Paredes, J.M. et al. 2000, Sci, 288, 2340
[583] Paredes, J.M. et al. 2002, A&A, 393, L99
[584] Parkes, G.E. et al. 1978, MNRAS, 184, 73P
[585] Parkes, G.E. et al. 1980, MNRAS, 190, 537
[586] Parkes, G.E. et al. 1980, MNRAS, 191, 547
[587] Parmar, A.N. et al. 1989, ApJ, 338, 359 & 373
[588] Parmar, A.N. et al. 1993, A&A, 275, 227
[589] Parmar, A.N. et al. 2000, A&A, 360, L31
[590] Parsignault, D.R. et al. 1972, Nat. Phys. Sci. 239, 123
[591] Patel, S.K. et al. 2004, ApJ, 602, L45
[592] Paul, B. & Rao, A.R. 1998, A&A, 337, 815
[593] Pavlinsky, M.N. et al. 1992, SvAL, 18, 88
[594] Payne, B.J. & Coe, M.J. 1987, MNRAS, 225, 985
[595] Pellizza, L.J. et al. 2005, A&A, submitted
[596] Penny, A.J. et al. 1973, MNRAS, 163, 7P
[597] Percy, J. et al. 1981, AJ, 86, 53
[598] Perryman, M.A.C. et al. 1997, A&A, 323, L49
[599] Phlips, B.F. et al. 1996, ApJ, 465, 907
[600] Pietsch, W. et al. 1980, ApJ, 237, 964
[601] Pietsch, W. et al. 1986, A&A, 163, 93
[602] Pigulski, A. et al. 2001, A&A, 376, 144
[603] Piraino, S. et al. 2000, A&A, 357, 501
[604] Polcaro, V.F. et al. 1990, A&A, 231, 354
[605] Polcaro, V.F. et al. 1993, A&A, 273, L49
[606] Pooley, G. 2004, ATel 276
[607] Pooley, G. G. et al. 1999, MNRAS, 302, L1
[608] Popper, D. 1950, ApJ, 111, 495
[609] Pottschmidt, K. et al. 2000, A&A, 357, L17
[610] Pottschmidt, K. et al. 2003, A&A, 407, 1039
[611] Pounds, K.A. et al. 1975, MNRAS, 172, 473
[612] Powell, C. R. et al. 2002, ATel 93
[613] Predehl, P. et al. 2000, A&A, 357, L25
[614] Priedhorsky, W.C. & Holt, S.S. 1987, SSRv, 45, 291
[615] Priedhorsky, W.C. & Holt, S.S. 1987, ApJ, 312, 743
[616] Priedhorsky, W.C. & Terrell, J. 1983, Nat, 303, 681
[617] Priedhorsky, W.C. & Terrell, J. 1983, ApJ, 273, 709
[618] Priedhorsky, W. & Terrell, J. 1984, ApJ, 280, 661
[619] Priedhorsky, W. & Terrell, J. 1986, ApJ, 301, 886
[620] Priedhorsky, W. et al. 1983, ApJ, 270, 233
[621] Punsly, B. 1999, ApJ, 519, 336
[622] Qu, J.L. et al. 2005, ApJ, 629, L33
[623] Quaintrell, H. et al. 2003, A&A, 401, 313
[624] Rappaport, S.A. et al. 1971, ApJ, 168, L17
[625] Rappaport, S.A. et al. 1978, ApJ, 224, L1
[626] Raubenheimer, B.C. 1990, A&A, 234, 172
[627] Raubenheimer, B.C. & Ogelman, H. 1990, A&A, 230, 73
[628] Ray, P.S. & Chakrabarty, D. 2002, ApJ, 581, 1293
[629] Ray, P.S. et al. 1997, ApJ, 491, 381
[630] Reid, C.A. et al. 1980, AJ, 85, 1062
[631] Reig, P. & Roche, P. 1999, MNRAS, 306, 95
[632] Reig, P. & Roche, P. 1999, MNRAS, 306, 100
[633] Reig, P. et al. 1996, A&A, 311, 879
[634] Reig, P. et al. 1997, A&A, 322, 183
[635] Reig, P. et al. 1998, MNRAS, 301, 42
[636] Reig, P. et al. 2000, MNRAS, 317, 205
[637] Reig, P. et al. 2001, A&A, 367, 266
[638] Reig, P. et al. 2003, A&A, 405, 285
[639] Reig, P. et al, 2004, A&A, 421, 673
[640] Reig, P. et al. 2005, A&A, 440, 637
[641] Reig, P. et al. 2005, A&A, 440, 1079
[642] Remillard, R.A. & Canizares, C.R. 1984, ApJ, 278, 761
[643] Remillard, R. & Levine, A. 1998, IAU Circ. 6826
[644] Remillard, R. & Smith, D. 2002, ATel 88
[645] Revnivtsev, M.G. 2003, AstL, 29, 644
[646] Revnivtsev, M. et al. 2002, A&A, 391, 1013
[647] Revnivtsev, M. et al. 2004, A&A, 4324, L5
[648] Reynolds, A.P. et al. 1992, MNRAS, 256, 631
[649] Reynolds, A.P. et al. 1999, A&A, 349, 873
[650] Rho, J. et al. 2004, HEAD, 8.1730
[651] Ribó, M. et al. 1999, A&A, 347, 518
[652] Ribó, M. et al. 1999, NewAR, 43, 545
[653] Ribó, M. et al. 2002, A&A, 384, 954
[654] Ribó, M. et al. 2006, A&A, 449, 687
[655] Richer, G. et al. 1976, ApJ, 204, L73
[656] Rickard, J.J. 1974, ApJ, 189, L113
[657] Ricke, G.H. & Lebofsky, M.J. 1985, ApJ, 288. 618
[658] Ricker, G.R. et al. 1973, ApJ, 184, 237
[659] Ricketts, M.J. et al. 1981, SSRv, 30, 399
[660] Robba, N.R. & Warwick, R.S. 1989, ApJ, 346, 469
[661] Robba, N.R. et al. 1992, ApJ, 401, 685
[662] Robba, N.R. et al. 2001, ApJ, 562, 950
[663] Robinson, E.L. et al. 2002, ApJ, 565, 1169
[664] Robinson, R.D. et al. 2002, ApJ, 575, 435
[665] Roche, P. et al. 1993, A&A, 270, 122
[666] Roche, P. et al. 1997, A&A, 322, 139
[667] Roche, P. et al. 1997, IAUC 6698
[668] Rodriguez, J. 2003, ATel 194
[669] Rodriguez, J.A. et al. 2003, A&A, 407, L41
[670] Rodriguez, J.A. et al. 2005, A&A, 432, 235
[671] Rodriguez, J.A. et al. 2005, MNRAS, accepted
[672] Romero, G. E. et al. 2001, A&A, 376, 599
[673] Romney, J.D. et al. 1987, ApJ, 321, 822
[674] Rosado, M. & Gonzalez, J. 1981, ReMAA, 5, 93
[675] Rose, L.A. et al. 1979, ApJ, 231, 919
[676] Rossi, C. et al. 1991, A&A, 249, L19
[677] Rothschild, R.E. & Soong, Y. 1987, ApJ, 315, 154
[678] Rothschild, R.E. et al. 1977, ApJ, 213, 818
[679] Rubin, B.C. et al. 1997, ApJ, 488, 413
[680] Rupen, M.P. et al. 2003, ATel 152
[681] Rupen, M.P. et al. 2003, ATel 184
[682] Sadakane, K. et al. 1985, ApJ, 288, 284
[683] Sakano, M. et al. 2000, PASJ, 52, 1141
[684] Sakano, M. et al. 2002, ApJS, 138, 19
[685] Salotti, L. et al. 1992, A&A, 253, 145
[686] Samus, N.N. et al. 1999, IAU Circ 7277
[687] Sanduleak, N. 1979, AJ, 84, 1319
[688] Saraswat, P. & Apparao, K.M.V. 1992, ApJ, 401, 678
[689] Sato, N. et al. 1986, ApJ, 304, 241
[690] Sato, N. et al. 1986, PASJ, 38, 731
[691] Schalinski, C.J. et al. 1995, ApJ, 447, 752
[692] Schreier, E. et al. 1972, ApJ, 172, L79
[693] Schreier, E.J. et al. 1976, ApJ, 204, 539
[694] Schulz, N. et al. 2002, ApJ, 564, L21
[695] Schulz, N.S. et al. 2002, ApJ, 565, 1141
[696] Schwartz, D.A. et al. 1980, AJ, 85, 549

[697] Schwartz, R.A. et al. 1991, ApJ, 376, 312
[698] Scott, D.M. et al. 1997, ApJ, 488, 831
[699] Sembay, S. et al. 1990, ApJ, 351, 675
[700] Seward, F.D. et al. 1976, MNRAS, 175, 39P
[701] Seward, F. et al. 1980, Nat, 287, 806
[702] Seward, F.D. et al. 1986, ApJ, 305, 814
[703] Seward, F.D. 1990, ApJS, 73, 781
[704] Sguera, V. et al. 2005, astro-ph/0509018
[705] Shafter, A.W. et al. 1980, ApJ, 240, 612
[706] Shrader, C.R. et al. 1999, ApJ, 512, 920
[707] Sidoli, L. et al. 2005, A&A, 429, L47
[708] Sidoli, L. et al. 2005, A&A, 440, 1033
[709] Singh, N.S. et al. 2002, A&A, 392, 161
[710] Slettebak, A. 1982, ApJS, 50, 55
[711] Smith, D.A. & Takeshima, T. 1998, IAU Circ. 7014
[712] Smith, D. et al. 1998, IAU Circ. 6855
[713] Smith, D.A. et al. 1999, IAU Circ 7253
[714] Smith, D.M. 2004, ATel 338
[715] Smith, D.M. et al. 1998, ApJ, 501, L181
[716] Smith, D.M. et al. 1998, ApJ, 503, 877
[717] Smith, M.A. et al. 2004, ApJ, 600, 972
[718] Smith, D.M. et al. 2006, ApJ, 638, 974
[719] Smithsonian Astrophysical Observatory Star Catalog (Washington D.C. 1966)
[720] Soffitta, P. et al. 1998, ApJ, 494, L203
[721] Soong, Y. & Swank, J. H. 1989, ttxa, symp, 617
[722] Spencer, R.E. 1984, MNRAS, 209, 869
[723] Stark, M.J.; Saia, M. 2003, ApJ, 587, L101
[724] Steele, I.A. & Clark, J.S. 2001, A&A, 371, 643
[725] Steele, I.A. et al. 1996, A&AS, 120, 213
[726] Steele, I.A. et al. 1998, MNRAS, 297, L5
[727] Steele, I.A. et al. 1999, A&AS, 137, 147
[728] Steiner, J.E. et al. 1984, ApJ, 280, 688
[729] Steiner, C. et al. 2005, ATel 672
[730] Stella, L. et al. 1985, ApJ, 288, L45
[731] Stevens, J.B. et al. 1997, MNRAS, 288, 988
[732] Stewart, G.C. et al. 1987, MNRAS, 228, 293
[733] Stickland, D. et al. 1997, MNRAS, 286, L21
[734] Stier, M. & Liller, W. 1976, ApJ, 206, 257
[735] Stirling, A.M. et al. 2001, MNRAS, 327, 1273
[736] Stirling, A.M. et al. 2002, MNRAS, 337, 657
[737] Stocke, J. et al. 1985, PASP, 97, 126
[738] Stollberg, M.T. et al. 1993, IAU Circ. 5836
[739] Stollberg, M.T. et al. 1999, ApJ, 512, 313
[740] Strom, R.G. et al. 1989, Nat, 337, 234
[741] Stubbings, R. 1999, IAU Circ 7253
[742] Sugizaki, M. et al. 2001, ApJS, 134, 77
[743] Sunyaev, R.A. et al. 2003, ATel 190
[744] Swank, J.H. & Markwardt, C.B. 2004, ATel 359
[745] Takeshima, T. & Chakrabarty, D. 1998, IAU Circ. 7016
[746] Takeshima, T. et al. 1991, PASJ, 43, L43
[747] Takeshima, T. et al. 1994, ApJ, 436, 871
[748] Takeshima, T. et al. 1998, IAU Circ. 6826
[749] Takeuchi, Y. et al. 1990, PASJ, 42, 287
[750] Tamura, K. et al. 1992, ApJ, 389, 676
[751] Tananbaum, H. et al. 1972, ApJ, 177, L5
[752] Tarasov, A.E. et al. 2003, A&A, 402, 237
[753] Tashiro, M. et al. 1991, MNRAS, 252, 156
[754] Tawara, Y. et al. 1989, PASJ, 41, 473
[755] Taylor, A.R. & Gregory, P.C. 1982, ApJ, 255, 210
[756] Taylor, A.R. & Gregory, P.C. 1984, ApJ, 283, 273
[757] Taylor, A.R. et al. 1992, ApJ, 395, 268
[758] Taylor, M. et al. 1995, AJ, 109, 396
[759] Telting, J.H. & Kaper, L. 1994, A&A, 284, 515
[760] Telting, J.H. et al. 1998, MNRAS, 296, 785
[761] Terrell, J. & Priedhorsky, W. 1984, ApJ, 285, L15
[762] Thackeray, A.D. 1970, MNRAS, 150, 215
[763] Thom, C. et al. 1986, A&A, 165, L13
[764] Thomas, R.M. et al. 1979, MNRAS, 188, 19
[765] Tigelaar, S. P. et al. 2004, MNRAS, 352, 1015
[766] Tjemkes, S.A. et al. 1986, A&A, 154, 77
[767] Tomsick, J.A. et al. 2003, IAUC, 8076
[768] Tomsick, J.A. et al. 2004, ATel 224
[769] Torii, K. et al. 1998, ApJ, 508, 854
[770] Torii, K. et al. 1999, ApJ, 523, L65
[771] Torrejon, J.M. & Orr, A. 2001, A&A, 377, 148
[772] Torrejon, J.M. et al. 2004, A&A, 423, 301
[773] Torres, D.F. et al. 2005, ApJ, 626, 1015
[774] Treves, A. et al. 1980, ApJ, 242, 1114
[775] Tsunemi, H. 1989, PASJ, 41, 453
[776] Tsunemi, H. & Kitamoto, S. 1988, ApJ, 334, L21
[777] Tueller, J. et al. 2005, ATel 591
[778] Tuohy, L.R. et al. 1988, in: "Physics of Neutron Stars and Black Holes", ed. Y. Tanaka, (Universal Academic Press, Tokyo), p. 93
[779] Ubertini, P. et al. 1991, ApJ, 366, 544
[780] Ubertini, P. et al. 1991, ApJ, 383, 263
[781] Ulmer, M.P. et al. 1972, ApJ, 178, L121
[782] Ulmer, M.P. et al. 1973, ApJ, 184, L117
[783] Unger, S.J. et al. 1992, MNRAS, 256, 725
[784] Unger, S.J. et al. 1998, A&A, 336, 960
[785] Unno, W. et al. 1990, PASJ, 42, 269
[786] Van der Klis, M. & Bonnet-Bidaud, J.M. 1984, A&A, 135, 155
[787] Van der Klis, M. & Bonnet-Bidaud, J.M. 1989, A&A, 214, 203
[788] Van der Klis, M. & Jansen, F. 1985, Nat, 313, 768
[789] van der Meer, A. et al. 2005, A&A, 432, 999
[790] van Kerkwijk, M.H. 1993, A&A, 276, L9
[791] Van Kerkwijk, M. et al. 1989, A&A, 209, 173
[792] Van Kerkwijk, M. et al. 1992, Nat, 355, 703
[793] Van Kerkwijk, M. et al. 1995, A&A, 303, 483
[794] van Loon, J.T. et al. 2001, A&A, 375, 498
[795] Van Oijen, J.G.J. 1989, A&A, 217, 115
[796] Van Paradijs, J. 1991, in: 'Neutron Stars, Theory and Observation', eds. J. Ventura & D. Pines (Kluwer), p. 289
[797] Van Paradijs, J. et al. 1977, A&AS, 30, 195
[798] Van Paradijs, J. et al. 1983, A&A, 124, 294
[799] Van Paradijs, J. et al. 1984, A&AS, 55, 7
[800] van Rensbergen, W. et al. 1996, A&A, 305, 825
[801] Vermeulen, R. et al. 1987, Nat, 328, 309
[802] Verrecchia, F. et al. 2002, A&A, 393, 983
[803] Verrecchia, F. et al. 2002, ATel 102
[804] Vestrand, W. et al. 1997, ApJ, 483, L49
[805] Vidal, N.V. 1973, ApJ, 186, L81
[806] Vieira, S.L.A. et al. 2003, AJ, 126, 2971
[807] Vijapurkar, J. & Drilling, J.S. 1993, ApJS, 89, 293
[808] Vikhlinin, A. et al. 1994, ApJ, 424, 395
[809] Vrtilek, S.D. et al. 1991, ApJS, 76, 1127
[810] Wackerling, L.R. 1970, Memoirs RAS 73, 153
[811] Wagner, R.M. et al. 1986, ApJ, 308, 152
[812] Wagner, R.M. et al. 1989, ApJ, 346, 971
[813] Walter, R. et al. 2003, A&A, 411, L427
[814] Wang, Z.R. et al. 1990, A&A, 240, 98
[815] Warwick, R.S. et al. 1981, MNRAS, 197, 865
[816] Waters, L.B.F.M. et al. 1988, A&A, 198, 200
[817] Waters, L.B.F.M. et al. 1989, A&A, 213, L19
[818] Waters, L.B.F.M. et al. 1989, A&A, 220, L1
[819] Watson, M.G. et al. 1981, MNRAS, 195, 197
[820] Watson, M.G. et al. 1982, MNRAS, 199, 915
[821] Watson, M.G. et al. 1983, ApJ, 273, 688
[822] Watson, M.G. et al. 1986, MNRAS, 222, 261
[823] Webster, B.L. & Murdin, P. 1971, Nat, 235, 37
[824] Wen, L. et al. 2000, ApJ, 532, 1119
[825] White, N.E. & Holt, S.S. 1982, ApJ, 257, 318
[826] White, N.E. & Swank, J.H. 1984, ApJ, 287, 856
[827] White, N.E. et al. 1976, MNRAS, 176, 201
[828] White, N.E. et al. 1978, Nat, 274, 664
[829] White, N.E. et al. 1980, ApJ, 239, 655
[830] White, N.E. et al. 1984, ApJ, 270, 711
[831] Whitlock, L. et al. 1989, ApJ, 338, 381
[832] Whitlock, L. et al. 1989, ApJ, 344, 371
[833] Wickramasinghe, D.T. et al. 1974, ApJ, 188, 167
[834] Wijnands, R. & van der Klis, M. 2000, ApJ, 528, L93
[835] Wilms, J. et al. 2005, astro-ph/0510193
[836] Wilson, C.A. et al. 1997, ApJ, 479, 388
[837] Wilson, C.A. et al. 1998, ApJ, 499, 820
[838] Wilson, C.A. et al. 1999, ApJ, 511, 367
[839] Wilson, C.A. et al. 2002, ApJ, 565, 1150
[840] Wilson, C.A. et al. 2002, ApJ, 570, 287
[841] Wilson, C.A. et al. 2003, ApJ, 584, 996
[842] Wilson, C.A. et al. 2003, ApJ, 596, 1220
[843] Wilson, C.A. et al. 2005, ApJ, 620, L99
[844] Wilson, C.A. et al. 2005, ApJ, 622, 1024
[845] Wilson R.B. et al. 1997, IAU Circ., 6586
[846] Wilson, R.W. 1963, AJ, 68, 181
[847] Wojdowski, P.S. et al. 2003, ApJ, 582, 959
[848] Wolff, S.C. & Morrisson, N.D. 1974, ApJ, 187, 69
[849] Wolinski, K.G. et al. 1996, ApJ, 457, 859

[850] Wood, K.S. et al. 1984, ApJS, 56, 507
[851] Wood, K.S. et al. 1991, ApJ, 379, 295
[852] Wu, C.C. et al. 1982, PASP, 94, 149
[853] Wu, C.C. et al. 1983, PASP, 95, 391
[854] Yamauchi, S. et al. 1995, PASJ, 47, 189
[855] Yang, S. et al. 1988, PASP, 100, 233
[856] Zamanov, R.K. et al. 1999, A&A, 351, 543
[857] Zdziarski, A.A. et al. 2002, ApJ, 578, 357
[858] Zhang, S. et al. 2005, ApJ, 630, L65
[859] Zhang, S.N. et al. 1996, A&AS, 120, 227
[860] Ziolkowski, J. 2005, MNRAS, 358, 851
[861] Zurita Heras, J.A. et al. 2005, A&A, astro-ph/0511115